

# Prevalence and Antibiotic Susceptibility of *Escherichia coli* Isolated from Early-Onset Sepsis in Shiraz, Iran

## ARTICLE INFO

### Article Type Original Research

#### Authors

Aliakbar Rezaei, PhD<sup>1</sup>  
Fatemeh Javanmardi, MSc<sup>1</sup>  
Neda Pirbonyeh, MSc<sup>2</sup>  
Hamid Reza Parsa, MD<sup>3</sup>  
Zahra Eskandari kootahi, MSc<sup>3</sup>  
Amir Emami PhD<sup>2\*</sup>

#### How to cite this article

Rezaei AA, Javanmardi F, Pirbonyeh N., Parsa HR., Eskandari kootahi Z., Emami A. Prevalence and Antibiotic Susceptibility of *Escherichia coli* Isolated from Early-Onset Sepsis in Shiraz, Iran. 2021;7(4): 305-310

<sup>1</sup> Vice-Chancellor for Treatment, health system research department, Shiraz University of Medical Sciences, Shiraz, Iran.

<sup>2</sup>Microbiology Department, Burn & Wound Healing Research Center, Shiraz University of Medical Sciences, Shiraz, Iran;

<sup>3</sup>Faculty Department of child and infant health, Shiraz University of Medical science, Shiraz, Iran.

#### \* Correspondence

Address: Microbiology Department, Burn & Wound Healing Research Center, Shiraz University of Medical Sciences, Shiraz, Iran.  
emami.microbia@gmail.com

#### Article History

Received: June 25 2021  
Accepted: September 25 ,2021  
Published: November 05 ,2021

## ABSTRACT

**Background:** Neonatal sepsis is a clinical syndrome in neonates, which is an uncommon but significant cause of morbidity and mortality in infants. The aim of this study was to evaluate the incidence of sepsis caused by *Escherichia coli* and its antibiotic resistance pattern as well as to assess the potential risk factors in neonates and maternal characteristics in Shiraz.

**Material & Method:** This retrospective study was performed on infants with sepsis in the first three days of life during February 2019 to March 2021. Patients' information was obtained using their hospital records and a questionnaire. All statistical analyses were conducted using SPSS software Ver. 18.0. A *p*-value <.05 was considered as statistically significant

**Findings:** During this study, a total of 250 positive blood cultures were reported for infants less than 3 days old. Of these, 21(8.4%) *E. coli* strains were isolated from 14 preterm and 7 term neonates. In all patients, the most effective antibiotic was meropenem, and the highest resistance was observed to cefoxitin.

**Conclusion:** Base on the present study results, *E. coli* is the most prevalent Gram-negative bacterium isolated in Shiraz. Premature birth and very low weight are the most important risk factors for developing early-onset sepsis.

**Keywords:** Early-onset sepsis, *E. coli*, Antibiotic susceptibility.

## CITATION LINKS

- [1] Martin RJ, Fanaroff AA, Walsh ... [2] Rennie JM, Robertson NC. Textbook of neonatology. E... [3] Stoll BJ, Hansen NI, Higgins RD, Fanaroff AA, Duara S, Goldberg R, et al... [4] Rafati M, Farhadi R, Nemati-Hevelai E, Chabra A. Determination of... [5] Moore MR, Schrag SJ, Schuchat A. Effects of... [6] Oza S, Lawn JE, Hogan DR, Mathers C, Cousens SN. Neonatal cause-of-death estimates for... [7] Bizzarro MJ, Dembry L-M, Baltimore RS, Gallagher PG. Changing patterns in neonatal *Escherichia coli*... [8] Jefferies AL. Management of term infants at increased risk for early-onset bacterial sepsis. *Paediatr Child Health*. 2017;22(4):223-8... [9] Polin RA. Management of neonates with suspected or proven early-onset bacterial sepsis. *Pediatrics*. 2012;129(5):1006-15... [10] Chen IL, Huang HC, Wu CT, Ou-Yang MC, Chung MY, Chen CC, et al... [11] Pazzi D, Klein I, Baker C. Bacterial sepsis and meningitis. In: Remington JS, editor. *Infectious disease of the fetus and newborn infant*. 6th ed. Philadelphia, PA: Saunders; 2006, 248-95... [12] Ehman D, Samedi VM, Kalaniti K, Daspal S. Neonatal *Escherichia coli* infection in twins: Clinical spectrum and management dilemma. *BMJ Case Rep*. 2021;14(1):e238470... [13] Schrag SJ, Farley MM, Petit S, Reingold A, Weston EJ, Pondo T, et al. Epidemiology of invasive early-onset neonatal sepsis, 2005 to 2014. *Pediatrics*. 2016;138(6):e20162013... [14] Tessema B, Lippmann N, Knüpfer M, Sack U, König B. Antibiotic resistance... [15] Tehrani FHE, Moradi M, Ghorbani N. Bacterial etiology and antibiotic resistance patterns in neonatal sepsis in Tehran during 2006-2014. *Iran J Pathol*. 2017;12(4):356-61... [16] Behmadi H, Borji A, Taghavi-Rad A, Soghandi L, Behmadi R. Prevalence and... [17] Hosseini MB, Abdoli Oskouei S, Heidari F, Sadat Sharif A, Salimi Z, Sharif SAA. Determination of the frequency of microbial agents and drug susceptibility... [18] Ghotaslou R, Soltani Ahari H, Ghorashi S. Study of the microbial etiologies and resistance pattern of... [19] Hammoud MS, Al-Taiar A, Al-Abdi SY, Bozaid H, Khan A... [20] Kung YH, Hsieh YF, Weng YH, Lien RI, Luo J, Wang Y, et al. Risk factors of... [21] Afsharpaiman S, Torkaman M, Saburi A, Farzaampur A, Amirjalali S, Kavehmanesh Z. Trends in incidence of neonatal sepsis and antibiotic susceptibility of... [22] Nikpay S, YadegarAzadi A, Mohamadi J, Soleymani A, Badfar G. Epidemiologic indicators of... [23] El-Amir MI, El-Feky MA, Elwafa DAA, Abd-Elmawgood EA. Rapid diagnosis of neonatal sepsis by PCR for detection of 16S rRNA gene, while blood culture and PCR results... [24] Aftab R, Iqbal I. Bacteriological agents of neonatal sepsis in NICU at Nishtar Hospital Multan. J... [25] Aurangzeb B, Hameed A. Neonatal... [26] Yadav NS, Sharma S, Chaudhary DK, Panthi P, Pokhrel P, Shrestha A, et al. Bacteriological profile of... [27] Herbst A, Källén K. Time between membrane rupture...

## Introduction

Neonatal sepsis is a clinical syndrome in neonates, which is an uncommon but significant cause of morbidity and mortality in infants, predominantly in developing countries; it is responsible for 30-50% of infant mortality in these populations<sup>[1,2]</sup>. Premature neonates are susceptible to nosocomial infections due to immaturity of the immune system and the need for numerous invasive procedures. Neonatal sepsis is divided into two categories based on the time of onset: early-onset sepsis (EOS) (in the first 72 hours of life) and late-onset sepsis (LOS) (after 72 hours)<sup>[3]</sup>.

Various factors are involved in the development of neonatal sepsis, including type of delivery, preterm birth, premature rupture of the fetal membrane (PROM), and maternal infection<sup>[4]</sup>. Bacterial infection is one of the prominent causes of neonatal death worldwide. In recent years, the most important cause of EOS has been group B *Streptococcus*. The use of intrapartum antibiotic therapy as prophylaxis has drastically reduced the prevalence of premature neonatal sepsis with this organism. However, the prevalence of infections with other antibiotic-resistant bacteria causing infections in preterm and very low birth weight (VLBW) infants has increased<sup>[5]</sup>. Among infants, one of the most common causes of EOS is *Escherichia coli*. This organism is responsible for about 24% of all sepsis cases. In the last two decades, *E. coli* has been the most common cause of EOS in premature infants as well as LOS. In hospitalized neonates, EOS caused by *E. coli* has a high mortality rate compared to Gram-positive organisms.<sup>[6, 7]</sup>

*E. coli* infection is a substantial cause of mortality and morbidity in neonates, particularly preterm ones. It is a major etiological agent of a variety of clinical manifestations, including sepsis, pneumonia, meningitis, skin and soft tissue infections<sup>[8]</sup>. Since the introduction of intrapartum antibiotic prophylaxis, the prevalence of *E. coli* has remained stable, but

its prevalence has increased in VLBW infants<sup>[7]</sup>. Empirical antibiotic therapy with ampicillin, gentamicin, or cefotaxime is essential for the successful treatment of neonatal sepsis. Therefore, treatment is commonly started before receiving blood culture results<sup>[9]</sup>. The prevalence and resistance of bacteria have increased over time due to the use of antibiotics in different countries. As a result, continuous epidemiological monitoring along with frequent local adjustment of susceptibility patterns to antimicrobial agents is essential to establish a rational treatment approach<sup>[10]</sup>.

**Objectives:** The aim of this study was to evaluate the incidence of sepsis caused by *E. coli* and its antibiotic resistance pattern as well as to assess the potential risk factors in neonates and maternal characteristics in Shiraz.

## Material and Method

This retrospective study was performed on infants less than 3 days old with sepsis in the teaching hospitals of Shiraz University of Medical Sciences during February 2019 to March 2021. The ethics code of this study (99-7819) was received from the Ethics Committee of Shiraz University of Medical Sciences. EOS is the sign and symptom of clinical sepsis in infants less than 3 days old. If the pathogen is isolated from the blood or CSF of patients, and the result of the first culture is positive, it is considered as the beginning of sepsis. Patients' information was obtained using their hospital records and a questionnaire. This information includes: sepsis type, age, gender, type of cultured pathogen, delivery method, antibiotic susceptibility, abortion, PROM, birth weight, and gestational age. Categorical variables were expressed as frequency and percentage (%), and continuous variables were expressed as mean. Continuous variables were analyzed by employing Student's t-test. Multiple logistic regression analysis was used to determine the most powerful risk factors such as demographic data, including age,

gender, etc. All statistical analyses were conducted using SPSS software, Ver. 18.0 (SPSS Inc., Chicago, IL, USA). A  $p$ -value  $<.05$  was considered as statistically significant.

### Findings

During this study, a total of 250 positive blood cultures were reported for infants less than 3 days old. After performing specific microbiological tests, Gram-positive bacteria were identified as the most common causes of infections. *Staphylococcus aureus* and coagulase-negative staphylococci with 156 (62.4%) and 53 (21.2%) isolates were the most prevalent Gram-positive bacteria in the studied samples. *E. coli* with 21 (8.4%) isolates was the most prevalent Gram-negative bacteria and the third most common cause of infection. Other isolated Gram-negative organisms included *Klebsiella pneumoniae* with 4 (1.6%) isolates, *Pseudomonas aeruginosa* with 3 (1.2%) isolates, *Acinetobacter baumannii* with 3

(1.2%) isolates, and *Enterobacter* species with 3 (1.2%) isolates, respectively. *E. coli* strains were isolated from 14 preterm and 7 term neonates. Antibiotic susceptibility of *E. coli* isolates was evaluated based on the principles of the CLSI table. In preterm infants, the highest sensitivity was observed to meropenem (86%), and the highest resistance was related to ceftazidime (52%). The highest resistance to ceftazidime (57%) was also reported in term neonates. In general, the most effective antibiotic against this organism was meropenem (80%). The results of antibiotic susceptibility testing are shown in Table 1. In this study, 12 (57%) infants with EOS were male, and 9 (43%) neonates were with VLBW. Also, seven mothers had a history of PROM. Among mothers with preterm infants, there were 7 (33%) cases of abortion and 4 (19%) cases of multiple abortion. Also, 4 (19%) neonates were born by natural delivery, and the other were delivered by caesarian section. Other information on the

**Table 1**) Antibiotic resistance of *E. coli* isolates in blood culture (early onset sepsis)

Antibiotics	Preterm		Term		Total	
	Susceptible N(%)	Resistance N(%)	Susceptible N(%)	Resistance N(%)	Susceptible N(%)	Resistance N(%)
Ciprofloxacin	8(57)	6(43)	5(71)	2(29)	13(62)	8(38)
Cotrimoxazole	11(79)	3(21)	5(71)	2(29)	16(76)	5(24)
Amikacin	6(43)	8(57)	4(57)	3(43)	10(48)	11(52)
Gentamicin	8(57)	6(43)	5(71)	2(29)	13(62)	8(38)
Ceftazidime	7(50)	7(50)	3(43)	4(57)	10(48)	11(52)
Imipenem	8(57)	6(43)	5(71)	2(29)	13(62)	8(38)
Meropenem	12(86)	2(14)	5(71)	2(29)	17(80)	4(20)
Cefepime	10(71)	4(29)	5(71)	2(29)	15(71)	6(29)
Ceftazidime	8(57)	6(43)	5(71)	2(29)	13(62)	8(38)
Cefixime	10(71)	4(29)	4(57)	3(43)	14(67)	7(33)

risk factors among preterm and term neonates is shown in Table 2.

### Discussion

Sepsis is one of the most important causes of infant mortality, especially in developing countries. It is estimated that about 2% of infants become infected before birth and about 10% in the first month of life [11]. *E. coli* infection in neonatal period is a major health concern. However, information on the prevalence, distribution, and clinical signs of *E. coli* infection in neonates is controversial and limited. In recent years, *E. coli* has surpassed

group B *Streptococcus* as the most important etiological agent of bacteremia in neonates in both preterm and term groups. Despite strict infection control programs in various countries, *E. coli* is the most common cause of severe infections such as meningitis with a mortality rate of over 40% [12, 13]. In this study, the most common Gram negative organism isolated was *E. coli*, which is consistent with other literature [14-16], but in contrast to other Iranian studies in which *P. aeruginosa* has been reported as the most prevalent organism [17, 18]. The difference in the results suggests that the microbial agents of newborn sepsis

**Table 2)** Risk factors for neonatal sepsis in teaching hospitals of Shiraz

Risk Factor		Preterm N(%)	Term N(%)	P-Value
Gender	Male	8(57)	4(57)	.24
	Female	6(43)	3(43)	
	Total	14(100)	7(100)	
PROM	Yes	5(36)	2(29)	.74
	No	9(64)	5(71)	
	Total	14(100)	7(100)	
VLBW	Yes	8(57)	1(14)	.03
	No	6(43)	6(86)	
	Total	14(100)	7(100)	
NVD	Yes	1(7)	3(43)	.049
	No	13(93)	4(57)	
	Total	14(100)	7(100)	
Abortion	Yes	4(28)	0(0)	.11
	No	10(72)	7(100)	
	Total	14(100)	7(100)	
Mother's age (year)		33.00±7.00	28.00±4.00	.25
Gestational age (week)		29.07±3.00	38.00±2.00	.04
Birth weight (gr)		1405±527	2410±717	.01

PROM: premature rupture of fetal membrane, VLBW: very low birth weight, NVD: normal vaginal delivery



may vary in different geographical locations and change over time. Therefore, various studies in different places and times are necessary in this regard. In the present study, the rate of sepsis in male neonates was higher than in female neonates, which is similar to other studies results in Iran, the Persian Gulf countries, and Taiwan [17-20]. Based on antibiotic susceptibility testing results in this study, the highest susceptibility in preterm neonates was to meropenem, cotrimoxazole, and cefepime, respectively, and the highest resistance was observed to amikacin. However, the highest resistance to cefoxitin was observed in term neonates as well as in all isolates. There are many reports about antibiotic resistance of routine EOS-related organisms [16, 21-23]. Aftab and Iqbal (2006) reported that resistance to gentamicin and cephalosporins was higher than to imipenem [24]. Another study by Aurangzeb and Hameed (2003) found that there was a higher resistance to routinely used antibiotics than to less commonly used antibiotics such as imipenem and ciprofloxacin [25]. In the present study, due to the routine use of gentamicin and amikacin, which are prescribed as prophylaxis, a higher rate of resistance to these antibiotics was observed. Unfortunately, despite the widespread use of ampicillin in hospitals, the degree of resistance to this antibiotic was not investigated in the present study.

Preterm birth and VLBW are the most important risk factors for EOS in infants [26]. In this study, 67% of patients with EOS were preterm neonates, and 57% had VLBW, while only 14% of them were term neonates. This suggests that preterm birth could be a double risk factor for VLBW newborns. In this study, only 20% of neonates were born by normal vaginal delivery. Almost one-third of our patients were exposed to PROM, which is consistent with other studies, concluding that the risk of neonatal sepsis increases with premature membrane rupture [27]. PROM and VLBW significantly increase the risk of early-onset sepsis, which is similar

to a study by Nikpay et al. (2018) in Ilam. The study of Al Amir et al. (2019) in Egypt is also similar to our study [23]. Limitations of this study include the small number of samples and the lack of evaluating antibiotic susceptibility to ampicillin, which is used as prophylaxis.

## Conclusion

Base on the present study results, *E. coli* is the most prevalent Gram-negative bacterium isolated in Shiraz. On the other hand, by examining the risk factors, it is concluded that premature birth and VLBW are the most important risk factors for developing EOS. The results of antibiotic susceptibility testing indicate that antibiotics should be reconsidered, and antibiograms should be performed.

## Acknowledgements

The authors of this article express their gratitude and thanks to the personnel of Hofez hospital.

**Ethical Permission:** approval of conduct study was obtain from Ethics committee of Shiraz university of medical science (IR.SUMS.RESEARCH.REC.1399.7819).

**Conflicts of Interests:** The authors declare that there is no conflict of interests with the organization that sponsored this research and publications arising from this research.

**Authors Contribution:** Conceptualization: AE, Data curation: HP, ZE; Formal analysis: FM; Funding acquisition: HP, ZK; Methodology: AAR, HP, ZE; Project administration: AE; Resources: AE; Software: FM; Supervision: AE; Validation: AAR; Visualization: AAR, FP, NP; Writing-original draft: AAR, NP; Writing-review and editing: AE, AAR.

**Fundings:** No found.

**Consent to participate:** Authors declare that all patients have signed the participation form.

## References

1. Martin RJ, Fanaroff AA, Walsh MC. Fanaroff and Martin's neonatal-perinatal medicine e-book: Diseases of the fetus and infant. Elsevier Health Sciences; 2014.

2. Rennie JM, Robertson NC. Textbook of neonatology. Elsevier. 1999.
3. Stoll BJ, Hansen NI, Higgins RD, Fanaroff AA, Duara S, Goldberg R, et al. Very low birth weight preterm infants with early onset neonatal sepsis: The predominance of gram-negative infections continues in the National Institute of Child Health and Human Development Neonatal Research Network, 2002–2003. *Pediatr Infect Dis J*. 2005;24(7):635-9.
4. Rafati M, Farhadi R, Nemati-Hevelai E, Chabra A. Determination of frequency and antibiotic resistance of common bacteria in late onset sepsis at the neonatal ward in Boooli-Sina Hospital of Sari, Iran. *J Babol Univ Med Sci*. 2014;16(6):64-71.
5. Moore MR, Schrag SJ, Schuchat A. Effects of intrapartum antimicrobial prophylaxis for prevention of group-B-streptococcal disease on the incidence and ecology of early-onset neonatal sepsis. *Lancet Infect Dis*. 2003;3(4):201-13.
6. Oza S, Lawn JE, Hogan DR, Mathers C, Cousens SN. Neonatal cause-of-death estimates for the early and late neonatal periods for 194 countries: 2000–2013. *Bull World Health Organ*. 2014;93(1):19-28.
7. Bizzarro MJ, Dembry L-M, Baltimore RS, Gallagher PG. Changing patterns in neonatal Escherichia coli sepsis and ampicillin resistance in the era of intrapartum antibiotic prophylaxis. *Pediatrics*. 2008;121(4):689-96.
8. Jefferies AL. Management of term infants at increased risk for early-onset bacterial sepsis. *Paediatr Child Health*. 2017;22(4):223-8.
9. Polin RA. Management of neonates with suspected or proven early-onset bacterial sepsis. *Pediatrics*. 2012;129(5):1006-15.
10. Chen IL, Huang HC, Wu CT, Ou-Yang MC, Chung MY, Chen CC, et al. Analysis of early-onset bloodstream infection due to Escherichia coli infection in premature babies. *Medicine*. 2017;96(32):e7748.
11. Pazzi D, Klein I, Baker C. Bacterial sepsis and meningitis. In: Remington JS, editor. *Infectious disease of the fetus and newborn infant*. 6<sup>th</sup> ed. Philadelphia, PA: Saunders; 2006, 248-95.
12. Ehman D, Samedi VM, Kalaniti K, Daspal S. Neonatal Escherichia coli infection in twins: Clinical spectrum and management dilemma. *BMJ Case Rep*. 2021;14(1):e238470.
13. Schrag SJ, Farley MM, Petit S, Reingold A, Weston EJ, Pondo T, et al. Epidemiology of invasive early-onset neonatal sepsis, 2005 to 2014. *Pediatrics*. 2016;138(6):e20162013.
14. Tessema B, Lippmann N, Knüpfer M, Sack U, König B. Antibiotic resistance patterns of bacterial isolates from neonatal sepsis patients at university hospital of Leipzig, Germany. *Antibiotics*. 2021;10(3):323.
15. Tehrani FHE, Moradi M, Ghorbani N. Bacterial etiology and antibiotic resistance patterns in neonatal sepsis in Tehran during 2006-2014. *Iran J Pathol*. 2017;12(4):356-61.
16. Behmadi H, Borji A, Taghavi-Rad A, Soghandi L, Behmadi R. Prevalence and antibiotic resistance of neonatal sepsis pathogens in Neyshabour, Iran. *Arch Pediatr Infect Dis*. 2016;4(2):e33818.
17. Hosseini MB, Abdoli Oskouei S, Heidari F, Sadat Sharif A, Salimi Z, Sharif SAA. Determination of the frequency of microbial agents and drug susceptibility pattern of the neonatal sepsis in the neonatal intensive care unit at Alzahra Hospital, Tabriz, Iran. *Iran J Neonatol*. 2019;10(4):33-40.
18. Ghotaslou R, Soltani Ahari H, Ghorashi S. Study of the microbial etiologies and resistance pattern of neonatal septicemia in Tabriz Pediatric Hospital. *J Ardabil Univ Med Sci*. 2007;7(2):155-9.
19. Hammoud MS, Al-Taiar A, Al-Abdi SY, Bozaid H, Khan A, AlMuhairi LM, et al. Late-onset neonatal sepsis in Arab states in the Gulf region: Two-year prospective study. *Int J Infect Dis*. 2017;55:125-30.
20. Kung YH, Hsieh YF, Weng YH, Lien RI, Luo J, Wang Y, et al. Risk factors of late-onset neonatal sepsis in Taiwan: A matched case-control study. *J Microbiol Immunol Infect*. 2016;49(3):430-5.
21. Afsharpaiman S, Torkaman M, Saburi A, Farzaampur A, Amirsalari S, Kavehmanesh Z. Trends in incidence of neonatal sepsis and antibiotic susceptibility of causative agents in two neonatal intensive care units in Tehran, IR Iran. *J Clin Neonatol*. 2012;1(3):124-30.
22. Nikpay S, YadegarAzadi A, Mohamadi J, Soleymani A, Badfar G. Epidemiologic indicators of neonatal sepsis in teaching hospitals of Ilam, Western Iran during (2012-2017). *Int J Pediatr*. 2018;6(7):7947-58.
23. El-Amir MI, El-Feky MA, Elwafa DAA, Abd-Elmawgood EA. Rapid diagnosis of neonatal sepsis by PCR for detection of 16S rRNA gene, while blood culture and PCR results were similar in E. coli-predominant EOS cases. *Infect Drug Resist*. 2019;12:2703-10.
24. Aftab R, Iqbal I. Bacteriological agents of neonatal sepsis in NICU at Nishtar Hospital Multan. *J Coll Physicians Surg Pak*. 2006;16(3):216-9.
25. Aurangzeb B, Hameed A. Neonatal sepsis in hospital-born babies: Bacterial isolates and antibiotic susceptibility patterns. *J Coll Physicians Surg Pak*. 2003;13(11):629-32.
26. Yadav NS, Sharma S, Chaudhary DK, Panthi P, Pokhrel P, Shrestha A, et al. Bacteriological profile of neonatal sepsis and antibiotic susceptibility pattern of isolates admitted at Kanti Children's Hospital, Kathmandu, Nepal. *BMC Res Notes*. 2018;11(1):1-6.
27. Herbst A, Källén K. Time between membrane rupture and delivery and septicemia in term neonates. *Obstet Gynecol*. 2007;110(3):612-8.