

Antibacterial Resistance and Virulence Potential of Avian Colibacillosis-Causing *Escherichia coli* Isolates

ARTICLE INFO

Article Type Original Article

Authors

Zohreh Pourhossein, MSc
Leila Asadpour, PhD*
Hadi Habibollahi, PhD
Seyedeh Tooba Shafighi, PhD

How to cite this article

Pourhossein Z, Asadpour L, Habibollahi H, Shafighi ST. Antibacterial Resistance and Virulence Potential of Avian Colibacillosis-Causing *Escherichia coli* Isolates. Infection Epidemiology and Microbiology. 2022;8(4): 307-315

Department of Biology, Rasht Branch,
Islamic azad University, Rasht, Iran.

* Correspondence

Department of Biology, Rasht
Branch, Islamic azad University,
Rasht, Iran.
Email: Asadpour@iaurasht.ac.ir

Article History

Received: April 08, 2022
Accepted: October 09, 2022
Published: December 19, 2022

ABSTRACT

Backgrounds: This study was done to evaluate the distribution of virulence-associated genes and antibiotic resistance in avian colibacillosis-causing *Escherichia coli* (*E. coli*) isolates.

Materials & Methods: In this study, 122 *E. coli* strains isolated from colibacillosis-suspected chickens in commercial broiler poultry farms (Guilan province, Iran) were examined for the presence of 12 virulence genes (*hlyF*, *iroN*, *iss*, *iutA*, *ompT*, *astA*, *tra*, *sfa-foc*, *papC*, *fimH*, *cvi/cva*, and *Tia*) using polymerase chain reaction (PCR). Antimicrobial susceptibility assessment was performed for the isolates using disc diffusion method against 19 antibiotics.

Findings: The *fimH*, *iut*, *tra*, *iss*, *iroN*, *hly*, and *ompT* genes were detected as the most prevalent genes among colibacillosis-causing isolates (more than 70%), while *sfa-foc* (S fimbriae and F1C fimbriae subunits) had the lowest frequency among colibacillosis-causing *E. coli* isolates (3.28%).

Conclusion: Virulence-associated genes were frequently detected in avian pathogenic *E. coli* strains. These findings could help better understand the pathogenicity potential of *E. coli* in poultry. Preventative measures are necessary to reduce food and environmental contamination with avian *E. coli* strains.

Keywords: *Escherichia coli*, Colibacillosis, Virulence potential, Poultry.

CITATION LINKS

[1] Guerra PR, Herrero-Fresno A, Pors SE, Ahmed S, ... [2] Ibrahim MS, Hussein AH, Eid AA, Lebdaq MA. Molecular ... [3] Li Y, Chen L, Wu X, Huo S. Molecular characterization ... [4] Sarowska J, Futoma-Koloch B, Jama-Kmieciak A, Frej-Madrzak M, ... [5] Cyoya PS, Rodrigues GR, Nishio EK, Medeiros LP, Koga VL, Pereira AP, et al. ... [6] Ewers C, Li G, Wilking H, Kieβling S, Alt K, Antáo EM, et al. Avian pathogenic, uropathogenic, and ... [7] Eibach D, Dekker D, Boahen KG, Akenten CW, ... [8] Peterson E, Kaur P. Antibiotic resistance ... [9] Ramos S, Silva V, Dapkevicius MD, Caniça M, ... [10] Badger S, Abraham S, Saputra S, Trott DJ, Turnidge J, Mitchell T, et al. Relative performance of ... [11] Chuppava B, Keller B, Meißner J, Kietzmann M, Visscher C. Effects of ... [12] da Silva GJ, Mendonça N. Association between ... [13] Momtaz H, Rahimi E, Moshkelani S. Molecular detection ... [14] Johnson JR, Stell AL. Extended virulence genotypes of ... [15] Johnson TJ, Wannemuehler Y, Doetkott C, Johnson SJ, ... [16] Yamamoto T, Echeverria P. Detection of the enteroaggregative *Escherichia coli* heat-stable enterotoxin ... [17] Ahmed AM, Shimamoto T, Shimamoto T. Molecular ... [18] Haghighi Khoshkhoo P, Pourtaghi H, Akbariazad G, Mokhayeri S. Virulence-associated gene profiles of ... [19] Stromberg ZR, Johnson JR, Fairbrother JM, Kilbourne J, Van Goor A, Curtiss 3rd R, et al. Evaluation of *Escherichia coli* ... [20] Eftekharian S, Ghorbanpoor M, Seyfi Abad Shapouri MR, Ghanbarpour R, Jafari RJ, Amani A. Frequency of selected ... [21] Asadi A, Salehi TZ, Jamshidian M, Ghanbarpour ... [22] Xu X, Sun Q, Zhao L. Virulence factors and ... [23] Rashki A, Abdi HA, Shookohi M. Prevalence of ... [24] Murase K, Martin P, Porcheron G, Houle S, Helloin E, Pénary M, et al. HlyF produced ... [25] Knöbl T, Micke Moreno A, Paixao R, Gomes TA, Vieira MA, ... [26] Johnson TJ, Siek KE, Johnson SJ, Nolan LK. DNA ... [27] Velhner M, Suvajdžić L, Todorović D, ... [28] Piva IC, Pereira AL, Ferraz LR, Silva RS, Vieira AC, Blanco JE, et al. ... [29] Sgariglia E, Mandolini NA, Napoleoni M, ... [30] Subedi M, Luitel H, Devkota B, Bhattarai RK, ... [31] Jahantigh M, Samadi K, Dizaji RE, Salari S. Antimicrobial resistance and prevalence of ... [32] Moniri R, Dastehgoli K. Fluoroquinolone-resistant ... [33] Essawy AA, Alotaibi NF. Energy transfer in polymeric film: Selective ... [34] Salehi TZ, Bonab SF. Antibiotics ... [35] Vanni M, Meucci V, Tognetti R, ... [36] Yazdanpour Z, Tadjrobehkar O, Shahkhal M. Significant ... [37] Karami N, Wold A, Adlerberth I. Antibiotic ...

Introduction

Escherichia coli (*E. coli*) is part of the intestinal flora of healthy broilers, but some strains of this bacterium, designated as avian pathogenic *E. coli* (APEC), are the causative agent of avian colibacillosis. Avian colibacillosis is mainly characterized by airsacculitis and respiratory problems in broiler chickens, leading to systemic infection. This disease is considered as the major cause of production reduction at all ages and economic losses in many poultry farms worldwide [1, 2].

Furthermore, APEC isolates are genetically similar to human extraintestinal pathogenic *E. coli* (ExPEC) strains, including uropathogenic *E. coli* (UPEC). They could potentially be transmitted to humans and cause extraintestinal infections, particularly through the food chain or direct contact with birds and their feces [1, 3, 4]. ExPEC has a complex phylogenetic structure and carries a wide range of virulence factors, including P fimbrial usher (*papC*, *papG*), extracellular adhesive fimbriae (*ecp*), salmochelin siderophore receptor (*iroN*, *fyuA*), aerobactin siderophore receptor (*iutA*), episomal outer membrane protein (*ompT*, *ompTp*, *tsh*), hemolysin (*hlyA*), and serum resistance-associated protein (*iss*). Virulence genes of these proteins may be present in pathogenicity islands (PAIs) or plasmids [4, 5]. However, the distribution of these genes could be different in the UPEC and APEC groups of ExPEC. While a set of virulence features including adhesion-related genes (*crl*, *fimC*, and *mat*) are detected in almost all APEC and UPEC strains regardless of their pathotype, other virulence factors such as *hra*, *papC*, and plasmid-related genes *cvi/cva*, *sitD* ep., *iucD*, *iss*, *traT*, and *iroN* are frequently detected in APEC. On the other hand, *iha*, *sfa/foc*, and *afa/draB* genes are more prevalent in UPEC than in APEC [6].

APEC strains could be a reservoir of virulence genes of ExPEC strains pathogenic to humans. Some human extraintestinal pathogenic *E. coli* strains have the *iss* gene in their genome

on the CoIV plasmid, which a huge virulence plasmid typical of avian pathogenic *E. coli* strains. This indicates that the exchange of plasmids and consequently these virulence genes is possible between APEC and UPEC strains [4].

Furthermore, the excessive use of antibiotics in animal husbandry for the treatment and prevention of infectious diseases or as growth promoters contributes to the emergence and dissemination of antibiotic-resistant bacteria [7, 8]. The most common transmission route of multidrug-resistant bacteria to humans is through the food chain [9]. Therefore, commensal and pathogenic *E. coli* strains containing various antimicrobial resistance genes entail a high risk of transmission of drug resistance to human microflora and other pathogenic bacteria [10, 11]. Antibiotic-resistant *E. coli* strains become even more alarming and life threatening if they are pathogenic. In addition, the acquisition of resistance determinants may represent an advantage for the survival of virulent *E. coli* strains. Virulence genetic determinants, if located on the same genetic platform as antimicrobial resistance genes, may be co-transferred under antimicrobial selective pressure, resulting in prolonged illness, greater risk of death, and higher chance of dissemination of resistant microorganisms [12].

Objectives: Surveillance of poultry-isolated *E. coli* is necessary for reducing their health risk and preventing infection. Although several studies have been performed on the virulence potential and antibiotic resistance of *E. coli* bacteria in different parts of Iran, data in Guilan province are very limited. Accordingly, this study aimed to investigate antibacterial resistance and the presence of virulence-associated genes in avian colibacillosis-causing *E. coli* strains in Guilan province, northern Iran.

Materials and Methods

Sampling and *E. coli* isolation: In this study,

122 *E. coli* strains were isolated from heart and liver samples collected from 20 to 45-day-old poultry with characteristic lesions consistent with colibacillosis in commercial broiler poultry farms in Guilan province (northern Iran) during 2020. The collected samples were transferred to MacConkey (MC) and Eosin Methylene Blue (EMB) agar media, and purification was performed according to standard microbiological and biochemical methods such as growth on triple sugar iron agar (TSI) and lysine iron agar (LIA). For oxidative/fermentative degradation of glucose, citrate utilization, urease production, indole production, glucose degradation (methyl red test), and motility tests were used [13].

Antimicrobial susceptibility testing: *E. coli* isolates were subjected to antimicrobial susceptibility testing to ceftriaxone (30 µg), cephalixin (30 µg), cefixime (5 µg), ceftazidime (30 µg), imipenem (10 µg), gentamicin (10 µg), amikacin (30 µg), neomycin sulfate (30 µg), fosbac (200 µg), fosfomicin (200 µg), doxycycline (10 µg), cotrimoxazole (10 µg), Chlortetracycline (10 µg), Ciprofloxacin (5 µg), Enrofloxacin (5 µg), Danofloxacin (10 µg), flumequine (30 µg), florfenicol (30 µg), and linco-spectin (10 µg), purchased from High Media-India, by disc diffusion method on Mueller-Hinton agar (Merck, Germany) according to the CLSIVET01 protocols (2018). *E. coli* ATCC 25922 was included for quality control, and breakpoints were in accordance with CLSI guidelines.

Polymerase chain reaction (PCR) screening for virulence-associated genes: One colony of each bacterial strain was purified, and bacterial genomic DNA was extracted using a GenElute bacterial genomic DNA kit (Sigma-Aldrich, St Louis, MO). The presence of 12 virulence-associated genes, including *hlyF*, *iroN*, *iss*, *iutA*, *ompT*, *astA*, *tra*, *sfa-foc*, *papC*, *fimH*, *cvi/cva*, and *Tia*, was assessed using PCR as described previously [6, 14-16]. PCR reactions (25 µL) were performed using 2 U of Taq

polymerase, 2.5 µL of 10X buffer, 10 mM dNTPs, 20 pmol of each primer, 4 mM MgCl₂, and 4 µL of the sample DNA. PCR products were analyzed using agarose gel electrophoresis (1%). Table 1 shows the sequences of oligonucleotide primers used in this study.

Findings

***E. coli* isolation and antimicrobial susceptibility testing:** In total, 122 *E. coli* isolates were obtained from colibacillosis-suspected chickens.

Table 2 shows the results of antibacterial susceptibility testing of *E. coli* isolates to common antibiotics. The highest resistance of colibacillosis-causing *E. coli* strains was related to enrofloxacin, chlortetracycline, and flumequin (more than 90%). No resistance was observed to imipenem and ceftriaxone among the tested isolates, and resistance to gentamicin, amikacin, cephalixin, cefixime, and ceftazidime was low.

Frequency of virulence-associated genes: The frequency of virulence-associated genes and their distribution among colibacillosis-causing *E. coli* isolates are shown in Table 3. Virulence-associated genes *fimH*, *iut*, *tra*, *iss*, *iroN*, *hly*, and *ompT* were identified as the most prevalent genes in colibacillosis-causing isolates (more than 70%). *Sfa-foc* (S fimbriae and F1C fimbriae subunits) had the lowest frequency among colibacillosis-causing *E. coli* isolates (3.28%).

The correlation between the presence of virulence factors and antibiotic resistance: Most virulence-associated genes were predominant in antibacterial-resistant strains. The *hly-f* gene was the most prevalent virulence gene among antibiotic-resistant isolates. Most *fimH* and *papC*-positive isolates were antibiotic-resistant. The *sfa-foc* gene was observed only in fluoroquinolone- and sulfonamide-resistant groups. Among the 12 virulence genes investigated, ten genes were detected in cephalixin- and cefixime-resistant isolates, and five genes including

Table 1) Nucleotide sequences of primers used in this study, amplicon size (bp), and specific PCR program for amplification of each gene

Target Gene	Function	Primer Sequence (5'-3')	Cycles Temp / Time (Seconds)				References		
			Initial Denaturation °c/min	Denaturation Extension	Annealing Extension	Final Extension °c/min			
<i>iroN-F</i>	Salmochelinsiderophore receptor gene	AATCCGGCAAAAGACGCAACCCGCT	94/4min	94/30S	63/30S	72/45S	72/10min	553	15
<i>iroN-R</i>		TTCGGGCAACCCCTGCTTTGACTTT							
<i>Hly-f-F</i>	Putative avian hemolysin	GGCCACAGTCGTTTAGGGTGTCTTACC	94/4min	94/30S	63/30S	72/45S	72/10min	450	15
<i>Hly-f-R</i>		GGGGGTTTAGGCATTCGGATACTCAG							
<i>iss-F</i>	Episomal increased serum survival gene	CAGCAACCCGAAACCACTTGATG	94/4min	94/30S	63/30S	72/45S	72/10min	323	15
<i>iss-R</i>		AGCATTGCCAGAGCGGCAGAA							
<i>OmpT-F</i>	Episomal outer membrane protease gene	TCATCCCGGAAGCCTCCCTCACTACTAT	94/4min	94/30S	63/30S	72/45S	72/10min	496	15
<i>OmpT-R</i>		TAGGGTTTGCTGCACCTGGCTTCTGATAC							
<i>ast A -F</i>	EAST1 (heat-stable cytotoxin associated with enteroaggregative E. coli)	TGCCATCAACACAGTATATCC	94/4min	94/30S	55/30S	72/40S	72/10min	116	16
<i>ast A -R</i>		TAGGATCCTCAGGTCCGGAGTGCAGGC							
<i>Tra-F</i>	Transfer protein	GTTGGTGGATGAGCACAG	94/4min	94/30S	58/30S	72/40S	72/10min	290	6
<i>Tra-R</i>		TAGTTACATCTTCCACCATCG							
<i>Sfa-foc-F</i>	S fimbriae (sialic acid-specific) and F1C fimbriae	CTCCGGAACTGGGTGCAT CTTAC	94/4min	94/30S	63.5/30S	72/45S	72/10min	410	14
<i>Sfa-foc-R</i>		CGGAGGAGTAATTACAAAACCTGGCA							
<i>Pap C-F</i>	Pilus associated with pyelonephritis	GTGGCAGTATGAGTAATGACCGTTA	94/4min	94/30S	63.5/30S	72/40S	72/10min	200	14
<i>Pap C-R</i>		ATATCCTTTCTCCAGGGATGCAATA							
<i>Fim H-F</i>	Type 1 fimbriae (D-mannose-specific adhesin)	TGCAGAACGGATAAGCCGTG	94/4min	94/30S	63.5/30S	72/45S	72/10min	508	14
<i>Fim H-R</i>		GCAGTCACCTGCCCTCCGGT							
<i>CVI/Cva-F</i>	Structural genes of colicin V operon (microcin ColV)	TCCAAGCGACCCCTTATAG	95/4min	94/30S	58/30S	72/50S	72/10min	679	6
<i>CVI/Cva-R</i>		CGCAGCATAGTTCATGCT							
<i>Iut-A-F</i>	Aerobactin siderophore receptor gene	GGCTGACATCATGGGAACCTGG	94/4min	94/30S	63/30S	72/40S	72/10min	302	15
<i>Iut-A-R</i>		5'- 5'- GGCTGGACATCATGGGAACCTGG							
<i>Tia-F</i>	Toxigenic invasion locus in ETEC strains	5'- 5AGCGCTTCCGTCAGGACTT	94/4min	94/30S	58/30S	72/45S	72/10min	512	6
<i>Tia-R</i>		ACCAGCATCCAGATAGCGAT							

Table 2) Antimicrobial resistance properties in AFEC isolates

Antibiotic	Sensitive Isolates (APEC Group) N=122 N(%)	Intermediate Isolates (APEC Group) N=122 N(%)	Resistant Isolates (APEC Group) N=122 N(%)
Amikacin	100 (82)	22 (18.03)	0(0)
Gentamicin	118(96.72)	3 (2.46)	1(0.82)
Neomycin Sulphate	37 (30.33)	50 (40.98)	35 (28.68)
Ciprofloxacin	18(14.75)	62 (50.82)	42(34.43)
Flumequin	2 (1.64)	7 (5.74)	113 (92.62)
Enrofloxacin	0 (0)	14 (11.47)	118 (96.72)
Danofloxacin	13 (10.66)	21 (17.21)	88 (72.13)
Co-trimaxazol	8 (6.58)	8 (6.58)	106 (86.88)
Chlortetracycline	2 (1.64)	3 (2.46)	117 (95.90)
Doxicyclin	26 (21.31)	12 (9.84)	84 (68.85)
Florphenicol	13 (10.66)	11 (9.02)	98 (80.33)
Lincospectin	103(84.43)	11 (9.1)	8 (6.56)
Imipenem	120(98.4)	2 (1.64)	0(0)
Cephalexine	89(73)	26 (21.31)	7(5.73)
Cefixime	21(17.21)	95 (78)	6(4.92)
Ceftazidime	119(97.54)	1 (0.82)	2(1.64)
Ceftriaxon	117 (96)	5 (4.1)	0(0)
Fosbac	98(80.32)	6 (4.92)	18(14.75)
Fosfomycin	80(65.57)	6 (4.92)	36(29.51)

hly-f, *iss*, *papC*, *cvi/cva*, and *iutA* were detected in ceftazidime-resistant isolates.

Table 3) Distribution of virulence associated genes in APEC strains

Gene	Frequency, N=122 N(%)
<i>iroN</i>	96(78.69)
<i>Hly-f</i>	72.95))89
<i>iss</i>	97 (79.51)
<i>ompT</i>	88 (72.13)
<i>astA</i>	63 (51.64)
<i>tra</i>	103 (84.43)
<i>Sfa-foc</i>	4(3.28)
<i>papC</i>	60(49.18)
<i>fimH</i>	122(100.00)
<i>CVI/Cva</i>	76 (62.29)
<i>Iut-A</i>	105 (86.06)
<i>Tia</i>	71 (58.2)

Discussion

APEC causes a variety of diseases and significant economic losses in the poultry industry worldwide. The expression of several virulence factors is responsible for the pathogenicity of APEC strains. In this study, the prevalence of 12 different virulence-associated genes was investigated in *E. coli* strains isolated from colibacillosis-suspected samples. Virulence-associated genes *fimH*, *iut*, *tra*, *iss*, *iut*, *iroN*, *hly*, and *ompT* were identified as the most prevalent genes in these isolates (more than 70%). Previous studies have found significant associations between these virulence genes and APEC strains [3,15,17]. In the present study, the prevalence rates of *iutA*, *hlyF*, *iroN*, *ompT*, and *iss* virulence genes among *E. coli* isolates were much higher than those reported in a

previous study on *E. coli* isolates collected from colibacillosis cases in broiler poultry farms in Alborz, Tehran, and Golestan provinces in Iran [18].

Studies have shown that the expression of surface adhesion factors and close contact between bacteria and the host cell wall increase the virulence of *E. coli* and help bacteria colonize infected host cells [4]. In the present study, the *fimH* gene had the highest frequency among the tested *E. coli* isolates. Consistent with this finding, Stromberg et al. (2017) identified the *fimH* gene in 90% of ExPEC isolates [19]. Another study showed that *fimH* was present in 61.7% of ExPEC isolates collected from chickens in southwestern Iran [20]. The prevalence of type 1 fimbrial adhesion gene (*fimH*) in APEC isolates in a study in southeastern Iran was 66.66%, and the *pap* gene was detected in 63.85% of these isolates [21].

The *ompT* gene promotes the formation of bacterial communities during APEC infection, and more than 80% of APEC isolates carry this gene, suggesting the essential role of this gene in APEC infection [22]. In this study, the prevalence of *ompT* was relatively high among the tested bacteria (72.13%). The *ompT* gene encodes an outer membrane protease (OmpT) that activates plasminogen to plasmin and plays an important role in protein degradation [23]. The *hlyF* gene is often found in APEC isolates and expressed during extraintestinal infection. This gene induces the formation of outer membrane vesicles (OMV) which are involved in the transmission of bacterial virulence factors and promote the pathological process in the infected host [24]. In this study, the *hlyF* gene was identified in 72.95% of APEC isolates. The frequency of *hlyF* in this study was lower than that previously reported by Li and colleagues (2015) [3]. However, the present study findings corroborated the results of other studies, which identified

the *hlyF* gene in most APEC strains and suggested it as a virulence marker associated with mammalian species and considered its presence in birds as an indicator of interspecies barrier transposition [24, 25].

The other virulence genes investigated, including *iroN* and *iutA*, encode outer membrane receptors for iron and siderophore aerobactin, respectively. In this study, these genes had high frequencies of 78.69 and 86.06%, respectively. Previous studies have shown that the prevalence of these genes is higher in APEC isolates compared with commensal *E. coli* [15, 26, 27]. The Tia protein contributes to invasive phenotypes and adhesion, and the *ast* gene encodes EAST1 protein, which is often compared with *E. coli* STa enterotoxin [28]. These genes were also detected in more than 50% of APEC strains in this study. Furthermore, in the present study, *sfa-foc* was identified in 3.28% of APEC isolates. S fimbria encoded by *sfa* operon is a common virulence factor among APEC and ExPEC strains. *E. coli sfa+* strains are very important since S fimbriae is associated with the pathogenesis of urinary infections, meningitis, and septicemia in human patients [25]. The low prevalence of *sfa-foc* in the present study was consistent with the findings of another study on *E. coli* isolates collected from pathological conditions of broiler chickens in poultry slaughterhouses in southeastern Iran [21].

Also, *cvi/cva*, as structural genes of colicin V operon (microcin ColV), was detected in more than 62.29% of APEC isolates. The frequency of this virulence gene was comparable to the literature. In a recently published study in Italy, the *cvi/cva* gene was detected in 16 (31%) out of 51 *E. coli* isolates [29]; also, Subedi et al. (2018) reported a prevalence of 57.8% for the *cvi/cva* virulence gene in APEC isolates collected from broiler chickens in Nepal [30].

Based on the results of antimicrobial susceptibility testing in the present study, *E. coli* isolates showed high drug resistance to at least two categories (quinolones and tetracycline) of antimicrobial agents. Among the 19 antimicrobial agents tested, APEC isolates showed the highest resistance to quinolones, such as flumequine (92.62%) and enrofloxacin (96.72%), as well as chlortetracycline (95.90%). Quinolones and tetracyclines are widely used to fight poultry pathogens in Iran [31, 32]. Flumequine is a broad-spectrum veterinary antibiotic that is widely used for prevention and treatment worldwide. Therefore, there are several reports of bacterial resistance to these antibacterial agents from all over the world [32-35]. Also, the high resistance rate to tetracyclines could be explained by the common use of oxytetracycline as a feed additive in commercial broiler rearing. On the other hand, less common antibiotics such as third-generation cephalosporins were found to be highly effective against *E. coli* isolates. These findings confirm that the incidence of antimicrobial resistance in *E. coli* isolates depends on the frequency of antibiotic use as feed additives or for the treatment of diseases in poultry. In this study, APEC isolates carried many virulence factors. Some virulence-related genes were predominantly associated with antibiotic resistance, which could cause a serious public health problem. The *hly-f* gene was found to be the most prevalent virulence gene among antibiotic-resistant isolates. Yazdanpour and co-workers (2020) also reported *hlyA* as the most prevalent virulence gene among highly resistant community-acquired uropathogenic *E. coli* isolates [36].

Also, adhesin-encoding genes *fimH* and *papC* were abundantly detected in antibiotic-resistant isolates in this study. In accordance with this result, some studies have shown

that the *papC* gene is a predictor of antibiotic resistance in uropathogenic *E. coli* isolates [36, 37]. Co-spreading of virulence factors and resistance genes indicates the presence of both groups of genes on the same transferable elements and horizontal gene transfer between bacteria.

Conclusion

The prevalence of virulence-associated genes and antibacterial resistance was high in colibacillosis-causing *E. coli* strains. Preventative measures are necessary to reduce food and environmental contamination with avian *E. coli* strains.

Acknowledgement

We gratefully acknowledge the support of Islamic Azad University, Rasht Branch, Rasht, Iran.

Conflicts of interests: The authors declare that there is no conflict of interest.

Ethical permissions: All applicable international, national, and/or institutional guidelines for animal care and use were followed.

Authors' contributions: Zohreh Pourhossein (first author): original researcher/ introduction author/ methodologist/ statistical analyst / discussion author; Leila Asadpour (second author): original researcher/ introduction author/ methodologist/ discussion author; Hadi Habibollahi (third author): original researcher/ data analyst; Seyedeh Tooba Shafighi (forth author): original researcher/ methodologist.

Fundings: None.

Consent to participate: This article does not contain any studies with human participants, and all procedures performed in this study were in accordance with animal care and use guidelines.

References

1. Guerra PR, Herrero-Fresno A, Pors SE, Ahmed S, Wang D, Thøfner I, et al. The membrane transporter PotE is required for virulence in

- avian pathogenic *Escherichia coli* (APEC). *Vet Microbiol.* 2018;216:38-44.
2. Ibrahim MS, Hussein AH, Eid AA, Lebdah MA. Molecular characterization of *Escherichia coli* strains causing respiratory signs in broiler chickens in Egypt. *Zag Vet J.* 2019;47(2):168-82.
 3. Li Y, Chen L, Wu X, Huo S. Molecular characterization of multidrug-resistant avian pathogenic *Escherichia coli* isolated from septicemic broilers. *Poult Sci.* 2015;94(4):601-11.
 4. Sarowska J, Futoma-Koloch B, Jama-Kmiecik A, Frej-Madrzak M, Ksiazczyk M, Bugla-Ploskonska G, et al. Virulence factors, prevalence, and potential transmission of extraintestinal pathogenic *Escherichia coli* isolated from different sources: Recent reports. *Gut Path.* 2019;11(1):1-6.
 5. Cyoia PS, Rodrigues GR, Nishio EK, Medeiros LP, Koga VL, Pereira AP, et al. Presence of virulence genes and pathogenicity islands in extraintestinal pathogenic *Escherichia coli* isolates from Brazil. *J Infect Dev Ctries.* 2015;9(10):1068-75.
 6. Ewers C, Li G, Wilking H, Kießling S, Alt K, Antão EM, et al. Avian pathogenic, uropathogenic, and newborn meningitis-causing *Escherichia coli*: How closely related are they? *Int J Med Microbiol.* 2007;297(3):163-76.
 7. Eibach D, Dekker D, Boahen KG, Akenten CW, Sarpong N, Campos CB, et al. Extended-spectrum beta-lactamase-producing *Escherichia coli* and *Klebsiella pneumoniae* in local and imported poultry meat in Ghana. *Vet Microbiol.* 2018;217:7-12.
 8. Peterson E, Kaur P. Antibiotic resistance mechanisms in bacteria: Relationships between resistance determinants of antibiotic producers, environmental bacteria, and clinical pathogens. *Front Microbiol.* 2018;9:2928.
 9. Ramos S, Silva V, Dapkevicius MD, Caniça M, Tejedor-Junco MT, Igrejas G, et al. *Escherichia coli* as commensal and pathogenic bacteria among food-producing animals: Health implications of extended spectrum β -lactamase (ESBL) production. *Animals.* 2020;10(12):2239.
 10. Badger S, Abraham S, Saputra S, Trott DJ, Turnidge J, Mitchell T, et al. Relative performance of antimicrobial susceptibility assays on clinical *Escherichia coli* isolates from animals. *Vet Microbiol.* 2018;214:56-64.
 11. Chuppava B, Keller B, Meißner J, Kietzmann M, Visscher C. Effects of different types of flooring design on the development of antimicrobial resistance in commensal *Escherichia coli* in fattening turkeys. *Vet Microbiol.* 2018;217:18-24.
 12. da Silva GJ, Mendonça N. Association between antimicrobial resistance and virulence in *Escherichia coli*. *Virulence.* 2012;3(1):18-28.
 13. Momtaz H, Rahimi E, Moshkelani S. Molecular detection of antimicrobial resistance genes in *E. coli* isolated from slaughtered commercial chickens in Iran. *Vet Med.* 2012;57(4):193-7.
 14. Johnson JR, Stell AL. Extended virulence genotypes of *Escherichia coli* strains from patients with urosepsis in relation to phylogeny and host compromise. *J Infect Dis.* 2000;181(1):261-72.
 15. Johnson TJ, Wannemuehler Y, Doetkott C, Johnson SJ, Rosenberger SC, Nolan LK. Identification of minimal predictors of avian pathogenic *Escherichia coli* virulence for use as a rapid diagnostic tool. *J Clin Microbiol.* 2008;46(12):3987-96.
 16. Yamamoto T, Echeverria P. Detection of the enteroaggregative *Escherichia coli* heat-stable enterotoxin 1 gene sequences in enterotoxigenic *E. coli* strains pathogenic for humans. *Infect Immun.* 1996;64(4):1441-5.
 17. Ahmed AM, Shimamoto T, Shimamoto T. Molecular characterization of multidrug-resistant avian pathogenic *Escherichia coli* isolated from septicemic broilers. *Int J Med Microbiol.* 2013;303(8):475-83.
 18. Haghighi Khoshkhou P, Pourtaghi H, Akbariazad G, Mokhayeri S. Virulence-associated gene profiles of avian pathogenic *Escherichia coli* isolated from broilers with colibacillosis: A pilot study in Iran. *Int J Enteric Pathog.* 2019;7(1):4-8.
 19. Stromberg ZR, Johnson JR, Fairbrother JM, Kilbourne J, Van Goor A, Curtiss 3rd R, et al. Evaluation of *Escherichia coli* isolates from healthy chickens to determine their potential risk to poultry and human health. *PloS One.* 2017;12(7):e0180599.
 20. Eftekharian S, Ghorbanpoor M, Seyfi Abad Shapouri MR, Ghanbarpour R, Jafari RJ, Amani A. Frequency of selected virulence-associated genes in intestinal and extra-intestinal *Escherichia coli* isolates from chicken. *Iran J Vet Med.* 2016;10(2):91-6.
 21. Asadi A, Salehi TZ, Jamshidian M, Ghanbarpour R. ECOR phylotyping and determination of virulence genes in *Escherichia coli* isolates from pathological conditions of broiler chickens in poultry slaughter-houses of southeast of Iran. *Vet Res Forum.* 2018;9(3):211-6.
 22. Xu X, Sun Q, Zhao L. Virulence factors and antibiotic resistance of avian pathogenic *Escherichia coli* in eastern China. *J Vet Res.* 2019;63(3):317-20.
 23. Rashki A, Abdi HA, Shookohi M. Prevalence of genes encoding outer membrane virulence factors among fecal *Escherichia coli* isolates. *Int J Basic Sci Med.* 2017;2(1):52-7.
 24. Murase K, Martin P, Porcheron G, Houle S, Helloin E, Pénary M, et al. HlyF produced by extraintestinal pathogenic *Escherichia coli* is a virulence factor

- that regulates outer membrane vesicle biogenesis. *J Infect Dis.* 2016;213(5):856-65.
25. Knöbl T, Micke Moreno A, Paixao R, Gomes TA, Vieira MA, da Silva Leite D, et al. Prevalence of avian pathogenic *Escherichia coli* (APEC) clone harboring *sfa* gene in Brazil. *Sci World J.* 2012;2012:437342.
 26. Johnson TJ, Siek KE, Johnson SJ, Nolan LK. DNA sequence of a ColV plasmid and prevalence of selected plasmid-encoded virulence genes among avian *Escherichia coli* strains. *J Bacteriol.* 2006;188(2):745-58.
 27. Velhner M, Suvajdžić L, Todorović D, Milanov D, Kozoderović G. Avian pathogenic *Escherichia coli*: Diagnosis, virulence, and prevention. *Arch Vet Med.* 2018;11(2):21-31.
 28. Piva IC, Pereira AL, Ferraz LR, Silva RS, Vieira AC, Blanco JE, et al. Virulence markers of enteroaggregative *Escherichia coli* isolated from children and adults with diarrhea in Brasilia, Brazil. *J Clin Microbiol.* 2003;41(5):1827-32.
 29. Sgariglia E, Mandolini NA, Napoleoni M, Medici L, Fraticelli R, Conquista M, et al. Antibiotic resistance pattern and virulence genes in avian pathogenic *Escherichia coli* (APEC) from different breeding systems. *Vet Ital.* 2019;55(1):27-33.
 30. Subedi M, Luitel H, Devkota B, Bhattarai RK, Phuyal S, Panthi P, et al. Antibiotic resistance pattern and virulence genes content in avian pathogenic *Escherichia coli* (APEC) from broiler chickens in Chitwan, Nepal. *BMC Vet Res.* 2018;14(1):1-6.
 31. Jahantigh M, Samadi K, Dizaji RE, Salari S. Antimicrobial resistance and prevalence of tetracycline resistance genes in *Escherichia coli* isolated from lesions of colibacillosis in broiler chickens in Sistan, Iran. *BMC Vet Res.* 2020;16(1):1-6.
 32. Moniri R, Dastehgoli K. Fluoroquinolone-resistant *Escherichia coli* isolated from healthy broilers with previous exposure to fluoroquinolones: Is there a link? *Microb Ecol Health Dis.* 2005;17(2):69-74.
 33. Essawy AA, Alotaibi NF. Energy transfer in polymeric film: Selective photoluminescent approach applied in durable optical sensing of flumequine in pharmaceutical and veterinary formulations. *Microchem J.* 2020;157:105081.
 34. Salehi TZ, Bonab SF. Antibiotics susceptibility pattern of *Escherichia coli* strains isolated from chickens with colisepticemia in Tabriz province, Iran. *Int J Poult Sci.* 2006;5(7):677-84.
 35. Vanni M, Meucci V, Tognetti R, Cagnardi P, Montesissa C, Piccirillo A, et al. Fluoroquinolone resistance and molecular characterization of *gyrA* and *parC* quinolone resistance-determining regions in *Escherichia coli* isolated from poultry. *Poult Sci.* 2014;93(4):856-63.
 36. Yazdanpour Z, Tadjrobehkar O, Shahkhah M. Significant association between genes encoding virulence factors with antibiotic resistance and phylogenetic groups in community acquired uropathogenic *Escherichia coli* isolates. *BMC Microbiol.* 2020;20(1):1-9.
 37. Karami N, Wold A, Adlerberth I. Antibiotic resistance is linked to carriage of *papC* and *iutA* virulence genes and phylogenetic group D background in commensal and uropathogenic *Escherichia coli* from infants and young children. *Eur J Clin Microbiol Infect Dis.* 2017;36(4):721-9.