

Isolation and Identification of Endophytic Fungi of Licorice and Their Effect on Human Pathogenic Fungi

ARTICLE INFO

Article Type Original Article

Authors Melika Esfandyari, *MSc¹* Reza Habibipour, *PhD*^{1*} Mohsen Rajabi, *PhD*²

¹ Department of Microbiology, Hamedan Branch, Islamic Azad University, Hamedan, Iran
² Forest and Rangeland Research Department, Hamedan Agriculture and Natural Resources Research and Education Center, AREEO, Hamedan, Iran

A B S T R A C T

Background: Endophytes are microorganisms residing within plants, which have recently gained attention for producing bioactive compounds with pharmacological properties like antioxidant, anti-inflammatory, anticancer, and antibiotic effects. Licorice plants (*Glycyrrhiza glabra*) may serve as a source of these bioactive metabolites, particularly those with antifungal potential. This study aimed to evaluate the antifungal properties of endophytic fungi residing in licorice plants against human pathogenic fungi. **Materials & Methods:** Licorice plant samples were collected from three locations in Hamadan in October 2022. Endophytic fungi were collected from different parts of licorice plants, identified morphologically using microscopy, and confirmed through molecular methods by DNA extraction and amplification, with sequencing results compared to public databases. The isolated fungi were cultured on PDA (potato dextrose agar) medium, and crude extracts were obtained using ethyl acetate. The antifungal properties of crude extracts were investigated against four fungal pathogens including *Aspergillus niger, Candida glabrata, Cryptococcus neoformans*, and *Pseudallescheria boydii* using disc diffusion method and ketoconazole as a positive control.

Findings: A total of 21 endophytic fungi were identified, with *Fusarium* as the dominant genus. *F. oxysporum* and *F. redolens* were prevalent in root tissues, while *A. niger* dominated in stems and leaves. *A. niger* exhibited strong antifungal effects against *A. niger*, *C. neoformans*, and *P. boydii*.

Conclusion: Licorice plant roots hosted the highest number of endophytes, likely due to reduced exposure to pollution. *A. niger* and *F. oxysporum* demonstrated significant potential as biocontrol agents against pathogenic fungi, making them promising candidates for sustainable disease management in agriculture and human health.

Keywords: Endophyte, Licorice, Pathogenic fungi

CITATION LINKS

* Correspondence

Department of Microbiology, Hamedan Branch, Islamic Azad University, Hamedan, Iran, Iran. Email: Habibipour@iauh.ac.ir

How to cite this article

Esfandyari M., Habibipour R., Rajabi M. Isolation and Identification of Endophytic Fungi of Licorice and Their Effect on Human Pathogenic Fungi Infection Epidemiology and Microbiology. 2025;11(2): 105-113.

Article History Received: July 03, 2024 Accepted: April 05, 2025 Published: April 21, 2025 [1] Yenn T, Ibrahim D. Endophytic fungi from Syzygium... [2] Bal C, Baba H, Akata I, Sevindik M, Selamoglu Z, Akgul H. Biological... [3] Strobel G, Daisy B, Castillo U, Harper J. Natural products... [4] Sevindik M, Akgul H, Selamoglu Z, Braidy N. Antioxidant and... [5] Sevindik M, Ozdemir B, Bal C, Selamoglu Z. Bioactivity of EtOH... [6] Esimone CO, et al. The isolation... [7] Jain P, Sharma P. Probiotics and their... [8] Wilson D. Endophyte: The evolution of a term... [9] Adeyemi AI. Isolation and screening of... [10] Strobel G, Daisy B. Bioprospecting for microbial... [11] Berdy J. Bioactive microbial metabolites... [12] Damle M. Glycyrrhiza glabra (Liquorice)-A potent... [13] Mohseni R, Noorbakhsh F, Moazeni M, Nasrollahi Omran A, Rezaie S. Antitoxin characteristic... [14] Dorman HD, Deans SG. Antimicrobial agents... [15] Salahvarzi S, Shakib P, Pirhadi M, Alikord M, Jahed Khaniki G. Helicobacter... [16] Kusari S, Lamshöft M, Zühlke S, Spiteller M. An endophytic... [17] Bhakuni DS, Dhar ML, Dhar MM, Dhawan BN, Mehrotra BN. Screening of... [18] Devi N, Prabakaran J. Bioactive metabolites... [19] Chowdhury NS, et al. Identification... [20] Alzoreky NS, Nakahara K. Antibacterial activity... [21] Gouda S, Das G, Sen SK, Shin HS, Patra JK. Endophytes... [22] Tiwari P, Bae H. Endophytic fungi... [23] Gu G, et al. Culturable... [24] Rachmawaty R, et al. Molecular... [25] Schoffen RP, et al. Evaluation of Trichoderma... [26] Xiao Y, et al. Antifungal screening of... [27] Sharma A, Kaushik N, Sharma A, Marzouk T, Djébali N. Exploring the... [28] Moussa AY. The limitless endophytes... [29] García-Latorre C, Rodrigo S, Santamaria O. Effect of fungal... [30] Costa D, Tavares RM, Baptista P, Lino-Neto T. Cork oak... [31] El-Sayed AS, et al. Exploiting the... [32] Xu J, Qi X, Zheng X, Cui Y, Chang X, Gong G. First report of... [33] Glass NL, Donaldson GC. Development of ...

Copyright@ 2025, TMU Press. This open-access article is published under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License which permits Share (copy and redistribute the material in any medium or format) and Adapt (remix, transform, and build upon the material) under the Attribution-NonCommercial terms.

Introduction

Human reliance on natural sources to discover novel pharmaceutical agents to combat diseases is well-documented ^[1]. Efforts to find safer, chemical-free, and less toxic therapeutic alternatives have driven the exploration of innovative approaches to address health challenges. As a result, numerous medications such as antifungal, antibiotic, antiulcer, anti-inflammatory, antidiabetic, and antioxidant agents have been derived from a variety of sources like medicinal and edible mushrooms (higher fungi), medicinal plants, and microbial reservoirs such as endophytes, bacteria, and fungi ^[2-5].

In recent years, bioactive metabolites of both known and novel endophytes have revealed diverse pharmacological properties, including antioxidant, anti-inflammatory, anticancer, antiviral, and antibiotic effects ^[6]. Endophytes are microorganisms that live within plant tissues without causing disease, which have recently emerged as promising reservoirs of new bioactive compounds [7]. Endophytes dwelling within the internal tissues of plants without causing symptomatic infections have been observed to confer protective benefits to their host plants through ^[8, 9]. the production of bioactive substances Remarkably, a single plant could host multiple endophytes, contributing to the rich

microbial diversity of plant ecosystems ^{[10,} ^{11]}. Licorice (*Glycyrrhiza glabra*) is a widely studied medicinal plant in Iran^[12]. The medicinal and flavoring properties of licorice have led to its extensive use in the pharmaceutical and food industries ^[1 3]. Licorice roots contain various secondary metabolites, including chalcones, coumarins, saponins, and flavonoids, which have antioxidant properties and offer numerous health benefits ^[14, 15]. Studies have demonstrated potent antimicrobial effects of licorice against fungi and bacteria, such as Candida albicans, Aspergillus niger, and A. flavus ^[13]. Therefore, endophytes in licorice plants may harbor a wealth of biologically active metabolites. This study aimed to evaluate the antifungal properties of endophytic fungi residing in licorice plants against human pathogenic fungi.

Objectives: The primary objective of the present study was to investigate the antifungal activity of endophytic fungal isolates collected from different parts of licorice plants (root, stem, and leaves) against human pathogenic fungi.

Materials and Methods

Collection of plant materials and isolation of endophytes: Samples of licorice plants (*Glycyrrhiza glabra*) were gathered from three locations (Moradbeg, Imamza-



Figure 1) Variations of licorice plants isolated from three regions: (A) Moradbeg, (B) Imamzadeh, and (C) Mavshan

deh, and Mavshan) in Hamadan during October 2022. Licorice plants were identified by a taxonomist in Hamadan, Iran (Figure 1). A testimonial sample (DACB-47653) of each plant was preserved in the herbarium for future reference.

The bark and leaves of healthy fully-grown plants were randomly selected for use in this research. Plant samples were collected in sterile bags and transported to the laboratory, where they were processed within a few hours of collection. Plant parts were placed on a clean glass plate and cut with a sterilized blade. Endophytic fungi were collected from healthy, fresh plant samples according to a modified protocol based on previous studies ^[16, 17].

Morphological identification of endophytic fungi: To identify the collected fungal isolates, they were placed on culture slides, stained with a lactophenol cotton blue reagent, and evaluated under phase-contrast and bright-field microscopes. Morphological identification of endophytic fungi was performed by considering characteristics mentioned in standard identification manuals, including conidia size and coloration, sporulation and acervulus production, aerial mycelium, margin characteristics, surface texture, colony and medium color, hyphal structure, and growth pattern ^[18].

Molecular identification of endophytic fungi: DNA amplification and internal transcribed spacer (ITS) sequencing were used for molecular identification of endophytic fungi following a molecular biological protocol ^[19, 20]. A small segment of fungal hyphae (0.5–1.0 cm²) was isolated from the Petri dish used for fungal cultivation and lyophilized in a 2 mL Eppendorf tube (Eppendorf, Germany). Lyophilized fungal mycelia were powdered, and DNA was extracted. Fungal DNA was extracted using the DNeasy Plant Mini Kit (Qiagen, USA) based on the manufacturer's protocol. The procedures included cell lysis, RNA digestion, removal of precipitates and debris, and DNA shearing, precipitation, and purification. The isolated DNA samples were amplified using PCR (polymerase chain reaction) and the HotStarTag Master Mix Kit (Qiagen, USA). The primers ITS1 and ITS4 (Invitrogen, USA) were mixed with the Hot Star Tag Master Mix Kit and DNA template in a total volume of 50 µL. Table 1 shows the primers used in this research.

The initial step was performed for 5 min at 95 °C. Subsequently, 35 cycles of PCR were conducted, which included the following steps: initial denaturation for 30 s at 95 °C, primer annealing for one min at 55.5 °C, and DNA amplification by polymerase for one min at 72 °C. Finally, a terminal extension step was carried out for 10 min at 72 °C. After the reaction, PCR products were electrophoresed on a 1% agarose gel at 90 volts and analyzed using a gel documentation system. Amplified fungal DNA samples (PCR products) were submitted for sequencing, and the resulting sequences were compared to publicly accessible databases, including GenBank, by employing the BLAST algorithm. Endophytic fungal cultivation and extraction: Cultivation of isolated fungal strains was performed on Petri dishes using approximately 1 L of PDA (potato dextrose agar) medium for each isolate. The fungal strain was incubated at 28 °C for 21 days, and the culture medium was extracted twice with ethyl acetate to obtain the crude extract

Table 1)	Primers	used in	the	present study
----------	---------	---------	-----	---------------

Primer Name	Nucleotide Sequence 3' -5'	Annealing Temperature	Reference
ITS1 (F)	TCCGTAGGTGAACCTGCGG	55.5	[32]
ITS4 (R)	TCCTCCGCTTATTGATATGC	55.5	[33]

^[19]. The fungal extract was condensed into a solid residue by solvent evaporation using a rotary vacuum evaporator under reduced pressure, yielding ethyl acetate crude extracts ^[20].

Antifungal activity test: The antifungal properties of isolated fungi and licorice plant parts were investigated against four fungal pathogens using the disc diffusion technique with slight modifications, including *A. niger*, *C. glabrata, Cryptococcus neoformans*, and *Pseudallescheria boydii* ^[20].

A ketoconazole disc $(30 \ \mu g/disc)$ served as a positive control. Solvents acted as negative controls. The inhibition zones around the discs were measured following incubation at 37 °C for 18 to 24 hours for bacteria and at 28 °C for 48 to 96 hours for fungi. The sensitivity of pathogenic fungi was determined by measuring the inhibition zone diameters in millimeters.

Statistical analysis: The experimental data obtained from the antifungal activity tests were recorded as the inhibition zone diameters (in mm).

All tests were performed in triplicate, and the findings were expressed as mean \pm SD (standard deviation). ANOVA (oneway analysis of variance) test was used to analyze differences between groups using GraphPad Prism software Version 9.0 (GraphPad Software, San Diego, CA, USA). Post hoc comparisons were conducted by employing Tukey's test to identify substantial differences between groups. The significance level was set at p < .05.

Findings

Morphological identification of fungal isolates: Totally, 21 endophytic fungal strains were collected from the leaves and bark of licorice plants, each of which was assigned an internal strain code: MS1, MS2, MS3, MS4, KS4, KS5, KR1, KR2, KR3, KR4, KR5, DR1, DR2, DR3, DR4, DR5, MR1, MR2, MR3, MR4, and MR5.

Microscopic examination revealed that strains DR1, DR2, and DR3 exhibited columnar conidial heads and flask-shaped vesicles with phialides covering half to three-quarters of the vesicle. The conidia of these strains were globose with a finely uneven surface, which displayed a plain green color, indicating their classification within *Aspergillus* species (Figure 2).

Strains MR1, MR2, MR3, MR4, MS3, KR1, and KR3 showed thinly-scattered to plentiful cottony mycelia with pigmentation ranging from colorless to pale violet and white to violet.

These strains contained hyaline conidia or phialospores and exhibited two spore types: unicellular oblong microconidia and multicellular macroconidia with slightly twisted or curved pointed ends. The colonies of these strains were fluffy with cylindrical spores that were either septate or aseptate. Accordingly, strains MR1, MR2, MR3, MR4, MS3, KR1, and KR3 were found to belong to *Fusarium* species (Figure 2). Other fungi were differentiated based on their unique microscopic characteristics.

Molecular identification of fungal



Figure 2) Microscopic structures of endophytes: (A) Aspergillus, (B) Fusarium, (C) Aspergillus, (D) Fusarium

isolates: To confirm the identity of the isolated strains, DNA sequence analysis was performed using Mega BLAST software in the U.S. National Center for Biotechnology Information (NCBI) databases. Table 2 provides information on endophytic fungi collected from licorice plants.

Antifungal screening of endophytic fungi isolated from licorice: To assess antifungal activity, extracts obtained from endophytic fungi were screened at a concentration of 100 µg per disc. *A. niger* extract was effective against all tested pathogenic fungi. Analysis of variance and evaluation of the average effect of *A. niger* endophytic fungus revealed that this endophyte had similar effects on *neoformans*.pathogenic fungi *A. niger* and *C* as the control. The other two pathogenic fungi, *P. boydii* and *C. glabrata*, showed less sensitivity to this endophyte than the

Table 2) NCBI BLAST results of fungal isolates recovered from licorice

Isolate	Species	Isolate	Species	
MR1	Fusarium f.sp	DR4	Fusarium f.sp	
MR2	Fusarium oxysporum	DR5	Fusarium f.sp	
MR3	Fusarium f.sp	MS2	Aspergillus niger	
MR4	Fusarium f.sp	KR1	Fusarium redolens	
MR5	Fusarium oxysporum	KR2	Fusarium redolens	
MS1	Aspergillus niger	KR3	Fusarium redolens	
MS3	Aspergillus niger	KR4	Fusarium redolens	
MS4	Aspergillus niger	KR5	Fusarium redolens	
DR1	Fusarium f.sp	KS4	Fusarium f.sp	
DR2	Fusarium f.sp	VCF	Fusarium f.sp	
DR3	Fusarium oxysporum	K35		



Figure 3) Comparison of the average effect of endophytic fungi isolated from licorice plant on the studied human pathogenic fungi

	Endophytic Fungi				
Patnogenic Fungus	Fusarium oxysporum	Fusarium redolens	Fusarium f.sp	Aspergillus niger	
Aspergillus niger	45.1	39.5	73.5	100.0	
Pseudallescheria boydii	27.2	27.6	9.3	43.1	
Cryptococcus neoformans	31.0	28.1	69.1	100.0	
Candida glabrata	23.4	24.9	82.2	83.9	
Control	-	-	-	-	

Table 3) Variance analysis of the effect of fungi isolated from licorice on the studied human pathogenic fungi

control, with the least sensitivity observed in the pathogenic fungus *P. boydi. F. redolens* extract was also found to be effective against pathogenic fungi, the highest and lowest inhibitory effects of this fungus were against pathogenic fungi A. niger and C. glabrata, respectively. Also, F. oxysporum f. sp showed .similar effects on pathogenic fungi C neoformans, A. niger, and C. glabrata, with the greatest effect observed against C. glabrata pathogenic fungus. F. oxysporum extract showed the highest and lowest inhibitory activities against pathogenic fungi A. niger and *C. glabrata*, respectively. The antifungal screening results could be observed in Table 3 and Figure 3.

Discussion

The demand for discovering novel chemical entities to combat various diseases is constantly increasing due to the emergence of new threats such as coronavirus disease 2019 (COVID-19), acquired immunodeficiency syndrome (AIDS), cancer, etc. Moreover, the global concern over antibiotic resistance has further amplified this need. Since ancient times, plants have been used as a source of potential biologically active compounds to combat various diseases. Re-

searchers have recently turned their attention towards plant-associated endophytes as they offer metabolites with greater remedial potential than the plants themselves ^[21]. Endophytic fungal strains are remarkable microorganisms that reside within plant tissues and are recognized as excellent reservoirs of biologically active compounds ^[22]. Therefore, investigating the biological activities of natural products, whether medicinal plants or fungi, in order to find potential candidate molecules is highly recommended. In this study, endophytes such as F. oxysporum, F. redolens, F. oxysporum f. sp., and A. niger were identified within licorice plants. In contrast, a similar study conducted in China on licorice plant endophytes identified F. oxysporum, F. sp. sporium, and F. graminearum, highlighting the diversity of endophytic species depending on geographical locations and environmental conditions ^[23]. The diversity of endophytes in licorice is in line with the findings of another research by Rachmawaty et al. (2024), where *F. sororula*, *Diaporthe* sp., and Sarocladium zeae were isolated from corn plants and demonstrated significant antifungal activity against *F. oxysporum*. These findings reinforce the importance of exploring different plant species and regions to uncover a broad spectrum of endophytic fungi with potential antifungal properties [24].

The antifungal activities observed in this study underscore the significant potential of endophytic fungi as biocontrol agents. A. niger endophytes showed the highest inhibitory activity against A. niger pathogenic fungi, demonstrating their potent biocontrol capability. Similarly, previous studies have shown that endophytes such as Trichoderma koningii and T. atroviride isolated from cedar have substantial growth inhibitory effects on A. niger, with Penicillium viridicatum showing a 51% inhibition rate in culture ^[25]. These results support the present study findings and suggest that various endophytic fungi possess strong antifungal properties against *A. niger* and other pathogenic fungi. Moreover, in a study by Xiao et al. (2013), endophytes isolated from Ginkgo biloba, specifically Chaetomium globosum, inhibited pathogenic fungi like F. graminearum and *Sclerotinia sclerotiorum*, illustrating the presence and versatility of a wide range of fungi with antifungal capabilities across different plant hosts ^[26]. Furthermore, this study revealed that A. niger endophytes exhibited significant inhibitory effects against C. neoformans, aligning with a previous research on Shirin-ebian plant endophytes that also showed inhibitory activity against *C. neoformans*; the two-way cross-culture method used in this previous research was effective in demonstrating the antifungal capabilities of these endophytes, providing a robust approach for screening potential biocontrol agents ^[27]. These findings align with the findings of another study by Moussa (2024), where endophytic fungi were highlighted as an underexplored resource of novel antifungal compounds effective against multidrug-resistant pathogens such as C. auris, C. albicans, C. neoformans, and A.

fumigatus. The potential of endophytic fungi to produce unique antifungal metabolites suggests a promising avenue for developing new antifungal agents, particularly against drug-resistant strains ^[28].

Interestingly, *F. oxysporum* showed high inhibitory activities against *P. boydii* and *C. glabrata*, mirroring the results of studies on fungal endophytes of cypress and other plants like *Nepeta crispa*. The inhibitory effects of endophytes such as *Alternaria radicina* and *Paecilomyces maximus* CAYPB58 against these pathogenic fungi further validate the antifungal potential of endophytic fungi ^[29-31].

The present research emphasizes the significant role of the plant root as a reservoir of endophytic fungi, likely due to its favorable microenvironment with lower pollution and dust levels. The findings highlight the importance of considering environmental factors influencing endophytic colonization patterns within plants. Identifying endophytes such as A. niger and F. oxysporum f. sp. as promising biocontrol agents underscores their potential use in sustainable disease management strategies. These endophytes could be harnessed for applications in crop protection and human health, reducing reliance on chemical fungicides and addressing concerns about antibiotic resistance.

Conclusion

This study revealed that the root part of the plant had the highest abundance of endophytes, likely attributable to lower levels of pollution and dust compared plant other parts. This finding to underscores the importance of considering factors influencing microenvironmental endophytic colonization patterns within plants. Moreover, this investigation identified fungal endophytes such as A. niger and F. oxysporum f. sp as particularly

promising candidates for biological control and management of pathogenic fungi. These findings not only shed light on the ecological dynamics within plant tissues but also highlight the potential of harnessing endophytic fungi for sustainable disease management strategies in agriculture and beyond. Further research in this area is warranted to elucidate the mechanisms underlying the biocontrol properties of these endophytes and to explore their practical applications in crop protection and human health.

Acknowledgments

The authors would like to acknowledge the Vice-Chancellor of Hamedan Branch of Islamic Azad University for supporting this study.

Conflicts of interests: The authors declare that they have no conflicts of interest.

Ethical permissions: Ethical approval was obtained from the Ethics Committee of Islamic Azad University, Hamedan Branch, Iran (code: IR.IAU.H.REC.1401.013).

Fundings/Support: This study was funded by Islamic Azad University, Hamedan Branch, Hamedan, Iran (Project No. 9608235279).

Authors' contributions: R.H. designed the present study and analyzed the obtained data, and M.E. contributed to experimental studies and sample collections. All authors read and approved the final version of the article.

Consent to participate: Written informed consent for participation in this study was obtained from the participants and/or their legal guardian(s). All techniques used in this study were conducted according to relevant regulations and guidelines. The findings were reported in accordance with STROBE guidelines.

References

1. Yenn T, Ibrahim D. Endophytic fungi from Syzygium cumini (L.) skeels leaves and its

potential as antimicrobial agents. IOP Conf Ser Earth Environ Sci. 2019:364(1):012023.

- Bal C, Baba H, Akata I, Sevindik M, Selamoglu Z, Akgul H. Biological activities of wild poisonous mushroom Entoloma sinuatum (Bull.) P. Kumm (Boletales). KSU J Agric Nat. 2022;25(1):83-7.
- 3. Strobel G, Daisy B, Castillo U, Harper J. Natural products from endophytic microorganisms. J Nat Prod. 2004;67(2):257-68.
- Sevindik M, Akgul H, Selamoglu Z, Braidy N. Antioxidant and antigenotoxic potential of infundibulicybe geotropa mushroom collected from northwestern Turkey. Oxid Med Cell Longev. 2020;2020(1):5620484.
- 5. Sevindik M, Ozdemir B, Bal Ć, Selamoglu Z. Bioactivity of EtOH and MeOH extracts of Basidiomycetes mushroom (Stereum hirsutum) on atherosclerosis. Arch Razi Inst. 2021;76(1):87-94.
- 6. Esimone CO, Treasure UN, Abba CC, Maduabuchi EP, Nnamdi OA, Ebiye EC, et al. The isolation, identification, and antimicrobial activities of endophytic fungi from Azadirachta indica. Biol Pharm Sci. 2020;11(03):115-24.
- 7. Jain P, Sharma P. Probiotics and their efficacy in improving oral health: A review. J Appl Pharm Sci. 2012;2(1):151-63.
- 8. Wilson D. Endophyte: The evolution of a term, and clarification of its use and definition. Oikos. 1995;73(2):274-6.
- 9. Adeyemi AI. Isolation and screening of endophytic fungi from three plants used in traditional medicine in Nigeria for antimicrobial activity. Int J Green Pharm. 2015;9(1):58-62.
- Strobel G, Daisy B. Bioprospecting for microbial endophytes and their natural products. Microbiol Mol Biol Rev. 2003;67(4):491-502.
- 11. Berdy J. Bioactive microbial metabolites. J Antibiot. 2005;58(1):1-26.
- 12. Damle M. Glycyrrhiza glabra (Liquorice)-A potent medicinal herb. Int J Herb Med. 2014;2(2):132-6.
- 13. Mohseni R, Noorbakhsh F, Moazeni M, Nasrollahi Omran A, Rezaie S. Antitoxin characteristic of licorice extract: The inhibitory effect on aflatoxin production in a spergillus parasiticus. J Food Saf. 2014;34(2):119-25.
- 14. Dorman HD, Deans SG. Antimicrobial agents from plants: Antibacterial activity of plant volatile oils. J Appl Microbiol. 2000;88(2):308-16.

- 15. Salahvarzi S, Shakib P, Pirhadi M, Alikord M, Jahed Khaniki G. Helicobacter phytotherapy: Medicinal plants affecting H. pylori infection in Iran. Plant Biotechnol Persa. 2020;2(2):38-23.
- Kusari S, Lamshöft M, Zühlke S, Spiteller M. An endophytic fungus from Hypericum perforatum that produces hypericin. J Nat Prod. 2008;71(2):159-62.
- 17. Bhakuni DS, Dhar ML, Dhar MM, Dhawan BN, Mehrotra BN. Screening of Indian plants for biological activity. II. Indian J Exp Biol. 1969;7(4):250-62.
- 18. Devi N, Prabakaran J. Bioactive metabolites from an endophytic fungus Penicillium sp. isolated from Centella asiatica. Curr Res Environ Appl Mycol. 2014;4(1):34-43.
- 19. Chowdhury NS, Sohrab MH, Rony SR, Suriya S, Begum MN, Rana MS, et al. Identification and bioactive potential of endophytic fungi from Monochoria hastata (L.) Solms. Bangladesh J Bot. 2016;45(1):187-93.
- 20. Alzoreky NS, Nakahara K. Antibacterial activity of extracts from some edible plants commonly consumed in Asia. Int J Food Microbiol. 2003;80(3):223-30.
- 21. Gouda S, Das G, Sen SK, Shin HS, Patra JK. Endophytes: A treasure house of bioactive compounds of medicinal importance. Front Microbiol. 2016;7:1538.
- 22. Tiwari P, Bae H. Endophytic fungi: Key insights, emerging prospects, and challenges in natural product drug discovery. Microorganisms. 2022;10(2):360.
- 23. Gu G, Jia X, Wang W, Li P, Zhao S, Zhou Z, et al. Culturable endophytic fungi from Glycyrrhiza inflata distributed in Xinjiang, China with antifungal activity. Microbiol Res. 2021;12(4):829-39.
- 24. Rachmawaty R, Mukrimah UL, Pagarra H, Junda M, Munisa A, Ali A, et al. Molecular Identification of endophytic fungi from corn plant as antifungal agent against Fusarium oxysporum. Trop J Nat Prod Res. 2024;8(8):8121-6.

- 25. Schoffen RP, Silva Ribeiro A, Oliveira-Junior V, Polonio JC, Polli AD, Orlandelli RC, et al. Evaluation of Trichoderma atroviride endophytes with growth-promoting activities on tomato plants and antagonistic action on Fusarium oxysporum. Cienc Nat. 2020;42:47.
- 26. Xiao Y, Li HX, Li C, Wang JX, Li J, Wang MH, et al. Antifungal screening of endophytic fungi from Ginkgo biloba for discovery of potent anti-phytopathogenic fungicides. FEMS Microbiol Lett. 2013;339(2):130-6.
- 27. Sharma A, Kaushik N, Sharma A, Marzouk T, Djébali N. Exploring the potential of endophytes and their metabolites for biocontrol activity. 3 Biotech. 2022;12(10):277.
- 28. Moussa AY. The limitless endophytes: Their role as antifungal agents against top priority pathogens. Microb Cell Fact. 2024;23(1):161.
- 29. García-Latorre C, Rodrigo S, Santamaria O. Effect of fungal endophytes on plant growth and nutrient uptake in Trifolium subterraneum and Poa pratensis as affected by plant host specificity. Mycol Prog. 2021;20(9):1217-31.
- Costa D, Tavares RM, Baptista P, Lino-Neto T. Cork oak endophytic fungi as potential biocontrol agents against Biscogniauxia mediterranea and Diplodia corticola. J Fungi. 2020;6(4):287.
- 31. El-Sayed AS, El-Sayed MT, Rady AM, Zein N, Enan G, Shindia A, et al. Exploiting the biosynthetic potency of taxol from fungal endophytes of conifers plants; genome mining and metabolic manipulation. Molecules. 2020;25(13):3000.
- 32. Xu J, Qi X, Zheng X, Cui Y, Chang X, Gong G. First report of Corynespora leaf spot on sweet potato caused by Corynespora cassiicola in China. Plant Dis. 2016;100(10):2163-.
- 33. Glass NL, Donaldson GC. Development of primer sets designed for use with the PCR to amplify conserved genes from filamentous ascomycetes. Appl Environ Microbiol. 1995;61(4):1323-30.