

ISSN: 2588-4115; Infection Epidemiology and Microbiology. 2019;5(1):7-14

Evaluation of Antimicrobial Resistance and Immune Evasion Cluster Genes in Clinical Methicillin-Resistant Staphylococcus aureus (MRSA) Isolates from Khuzestan Province, Iran

ARTICLE INFO

Article Type Original Research

Authors Ariyarad S.¹ *MSc*,

Rezatofighi S. E.^{2,1*}*PhD*, Motamedi H.^{2,1} *PhD*

How to cite this article

Ariarad S, Rezatofighi SE, Motamedi H. Evaluation of Antimicrobial Resistance and Immune Evasion Cluster Genes in Clinical Methicilin-Resistant *Staphylococcus aureus* (MRSA) Isolates from Khuzestan Province, Iran. Infection Epidemiology and Microbiology. 2019;5(1):7-14

¹ Department of Biology, Faculty of Science, Shahid Chamran University of Ahvaz, Ahvaz, Iran ² Biotechnology and Biological Science Research Center, Shahid Chamran University of Ahvaz, Ahvaz, Iran

* Correspondence

Address: e.tofighi@yahoo. com; e.tofighi@scu.ac.ir Postal code: 6135743135, Tel and Fax: 00986113331045 Postal Code: 1316943551

Article History

Received: February 14 ,2019 Accepted: Marc 14 ,2019 ePublished: May 30 ,2019

ABSTRACT

Aims: Methicillin-resistant *Staphylococcus aureus* (MRSA) is recognized as an important health problem worldwide. To counteract the human innate immunity, *S. aureus* produces a number of immune evasion clusters (IEC) including staphylokinase (SAK), staphylococcal enterotoxin P (SEP), staphylococcal enterotoxin A (SEA), staphylococcal complement inhibitor (SCIN), and chemotaxis inhibitory protein (CHIP), encoded by *sak*, *sep*, *sea*, *scn*, and *chp* genes, respectively. These genes are carried by β-haemolysin-converting bacteriophages. The present study was conducted to determine the IEC phage types and antibiotic resistance patterns in 145 clinical MRSA isolates from Khuzestan Province, Iran.

Materials & Methods: All the isolates were investigated by disc diffusion method and PCR assay for *sak*, *sep*, *sea*, *scn*, and *chp* genes.

Findings: The assessment of antibiotic resistance showed the highest resistance rate towards penicillin (97.25%), followed by methicillin (95.8%), ceftazidime (81.4%), erythromycin (71.8%), clindamycin (61.4%), ciprofloxacin (60.7%), gentamycin (56%), imipenem (56.55%), and vancomycin (0%), respectively. Also, the frequency of IEC types was as follows: Type A, 4.8%; Type B, 9%; Type C, 13.1%; Type D, 12.4%; Type E, 27.6%; Type F, 1.4%; Type G, 0.7%; and Type H, 6.9%. On the other hand, 24.1% of the isolates showed no IEC type. **Conclusion**: The findings showed that IEC-carrying bacteriophages were highly prevalent among the MRSA strains, resulting in the adaptation and counteraction of bacteria with the human immune system. Therefore, understanding the role of IEC in bacteria virulence can improve our knowledge about the evolution, vaccination, and treatment of *S. aureus* infection.

Keywords: Immune evasion cluster; Methicillin-resistant *Staphylococcus aureus*; Phage typing; Antibiotic resistance

CITATION LINKS

[1] Baptistão LG, Silva NC, Bonsaglia EC, Rossi BF, Castilho IG, Fernandes Junior A, et al. Presence of immune evasion cluster and molecular ... [2] Liang BS, Huang YM, Chen YS, Dong H, Mai JL, Xie YQ, et al. Antimicrobial resistance and ... [3] Hau SJ, Sun J, Davies PR, Frana TS, Nicholson TL. Comparative prevalence of immune evasion ... [4] McCarthy AJ, Witney AA, Lindsay JA. Staphylococcus aureus temperate bacteriophage... [5] Foster TJ. Immune evasion by staphylococci... [6] Yu F, Li T, Huang X, Xie J, Xu Y, Tu J, et al. Virulence gene profiling and... [7] Jongerius I, von Köckritz-Blickwede M, Horsburgh MJ, Ruyken M, Nizet V, Rooijakkers SH. Staphylococcus... [8] van Wamel WJ, Rooijakkers SH, Ruyken M, van Kessel KP, van Strijp JA. The innate ... [9] Verkaik NJ, Benard M, Boelens HA, de Vogel CP, Nouwen JL, Verbrugh HA, et al. Immune evasion ... [10] Jongerius I, Köhl J, Pandey MK, Ruyken M, van Kessel KP, van Strijp JA, et al. Staphylococcal ... [11] Kraushaar B, Hammerl JA, Kienöl M, Heinig ML, Sperling N, Dinh Thanh M, et al. ... [12] de Jong NWM, Vrieling M, Garcia BL, Koop G, Brettmann M, Aerts PC, et al. Identification ... [13] Wayne PA. CLSI, Performance standards for... [14] Emaneini M, Bigverdi R, Kalantar D, Soroush S, Jabalameli F, Noorazar Khoshgnab B, et al. Distribution of... [15] Ahmadrajabi R, Layegh-Khavidaki S, Kalantar-Neyestanaki D, Fasihi Y. Molecular analysis of ... [16] Price LB, Stegger M, Hasman H, Aziz M, Larsen J, Andersen PS, et al. ... [17] Stefani S, Chung DR, Lindsay JA, Friedrich AW, Kearns AM, Westh H, et al. Meticillin-resistant Staphylococcus aureus (MRSA): global epidemiology and harmonisation of... [18] Hudson LO, Murphy CR, Spratt BG, Enright MC, Elkins K, Nguyen C, et al. Diversity of... [19] Dinges MM, Orwin PM, Schlievert PM. ... [20] You Y, Song L, Nonyane BAS, Price LB, Silbergeld EK. Genomic differences between nasal ... [21] Zadoks RN, van Leeuwen WB, Kreft D, Fox LK, Barkema HW, Schukken YH, et al. Comparison of ... [22] Goerke C, Wirtz C, Fluckiger U, Wolz C. Extensive phage... [23] Sadeghi J, Mansouri S. Molecular characterization and ... [24] Ohadian Moghadam S, Pourmand MR, Mahmoudi M, Sadighian H. ... [25] Saderi H, Emadi B, Owlia P. Phenotypic and genotypic study of ...

Copyright© 2019, TMU Press. This open-access article is published under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License which permits Share (copy and redistribute the material in any medium or format) and Adapt (remix, transform, and build upon the material) under the Attribution-NonCommercial terms.

Introduction

Staphylococcus aureus as an opportunistic pathogen can colonize the skin and nasopharynx of healthy individuals. This bacterium is responsible for various human diseases, including bacteremia, skin infections, toxic shock syndrome, necrotizing pneumonia, osteomyelitis, and endocarditis ^[1, 2, 3]. Evidence suggests that *S*. aureus has a remarkable potential to acquire antimicrobial resistance. The first case of resistance to methicillin was reported in 1961 ^[3]. *mecA* gene located on a mobile genetic element was found to be responsible for ... resistance; this gene is also the cause of resistance to other β -lactam antibiotics. Methicillin-resistant S. aureus (MRSA) infection is still a major global concern due to the difficulty in its treatment, particularly isolates acquire when the multidrug resistance (MDR) elements, especially against the vancomycin [4, 5]. In this regard, Yu and collogues reported that MRSA isolates harbor more virulence genes than other isolates ^[6].

Generally, the virulence factors of S. aureus include toxins, adhesions, immune evasion proteins, and pore-forming proteins ^[1, 2]. The first-line defense against the bacterial infections is innate immunity ^[5]. To evade this part of the immune system, S. aureus expresses some small proteins ^[7]. One of these human-specific immune modulators is the staphylococcal complement inhibitor (SCIN). SCIN is a C3 convertase inhibitor, which prevents the generation of C3b component and opsonophagocytosis of S. aureus by neutrophils ^[8, 9]. The chemotaxis inhibitory protein of S. aureus (CHIPS), as another immune-modulating protein, specifically blocks phagocyte activation by C5a through binding to C5a receptor (C5aR) and formylated peptide receptor (FPR)^[8, 10]. Staphylococcal enterotoxin A (SEA) is a superantigen, which can regulate

8

the operation of chemokine receptors on monocytes, such as CCR1, CCR2, and CCR5^[8, 9]. Another superantigen, which limits the host's innate immune response to S. aureus antigens, is staphylococcal enterotoxin P (SEP). SEP also prevents bacterial elimination ^[3]. SEA and SEP can poisoning; foodborne however, cause harboring these enterotoxins strains are often methicillin-sensitive ^[11]. On the other hand, staphylokinase (SAK) secreted by S. aureus modulates the innate immune system through anti-opsonic activities and defensins destruction [8]. SAK converts plasminogen to plasmin and causes the cleavage of important opsonic molecules, including IgG and C3b; therefore, phagocytosis of staphylococci by neutrophils is prevented ^[9]. SCIN, CHIPS, SEA, SEP, and, SAK are encoded by scn, chp, sea, sep, and sak genes, respectively to form an immune evasion cluster (IEC) [8]. So far, eight IEC variants have been identified, which are designated as A to H^[8,9]. scn gene is common among the all variants. Except for Variant H, other variants harbor a combinations of scn gene with one or more other genes, including chp, sak, sea, and sep genes ^[1]. IECs are carried by bacteriophages and integrated with the β -hemolysinencoding genes. Consequently, these bacteriophages are known as β-hemolysinconverting phages ^[9]. Evidence shows that these phages are found in more than 90% of the clinical *S. aureus* isolates from humans; however, not all genes are present on a given bacteriophage, and gene combinations are diverse ^[3, 12].

Objectives: The aim of this study was to characterize MRSA isolates, based on the antibiotic resistance pattern and β -hemolysin-converting phages. Our findings can help design therapeutic strategies and preventive measures against the infection transmission.

Materials and Methods

Sample collection and characterization: The isolates were collected from the clinical samples of patients, admitted to the affiliated hospitals of Ahvaz Jundishapur University of Medical Sciences from April 2015 to December 2015. The clinical samples were collected from wounds (N: 67), urine (N: 16), and blood samples (N: 23); respiratory infections (N: 19); abscesses (N: 9); and catheters (N: 11). Conventional biochemical methods were used for the detection of S. aureus isolates, including Gram staining, DNase, catalase, furazolidone (100 µg) susceptibility testing, bacitracin (0.04 units) resistance tests, slide and tube coagulase tests, and mannitol fermentation on mannitol salt agar.

Detection of MRSA isolates: According to the Clinical & Laboratory Standards Institute (CLSI) guidelines for the detection of MRSA isolates, resistance to cefoxitin (30 µg) was investigated on Mueller-Hinton Agar (MHA) ^[13]. Then the resistant isolates were confirmed as MRSA with respect to the presence of *mecA* gene using the PCR method, as described previously ^[14].

Antimicrobial susceptibility testing: The isolates susceptibility to antibiotics was examined by disk diffusion method according to the CLSI guidelines (13). The antibiotics disks included imipenem (30 μ g), clindamycin (2 μ g), gentamicin (10 μ g), methicillin (5 μ g), erythromycin (15 μ g), penicillin (10 units), ciprofloxacin (5 μ g), and ceftazidime (30 μ g). The vancomycin-resistant *S. aureus* (VRSA) isolates were screened by culturing on the brain-heart infusion (BHI) agar using 6 μ g/ml of vancomycin ^[15].

DNA extraction: The strains were cultured on the BHI broth overnight at 37 °C. Then chromosomal DNA was extracted using the Gram-positive bacteria DNA extraction kit (SinaClon; Iran) based on the manufacturer's instructions.

IEC profiling: The complement of IEC genes was determined by the PCR assay to

Ariyarad S. et al

investigate the phage types of the isolates. The primers and conditions used for the detection of *chp*, *sak*, *sea*, *sep*, and *scn* genes were previously described by van Wamel *et al.* (2006) (Table 1) ^[8]. The isolates typing scheme was from A to H according to the studies done by van Wamel *et al.* (2006) and Price *et al.* (2012) (Table 2) ^[8, 16].

Table 1) Names and sequences of primers used inthis study

Primer name	Sequence (5'→3')	References
sak	AGGCGATGACGCGAGTTAT GCGCTTGGATCTAATTCAAC	8
scn	AGCACAAGCTT GCCAACATCG TTAATATTTACTTTTTTAGTGC	8
chp	TTTACTTTTGAACCGTTTCCTAC CGTCCTGAATTCTTAGTATGCATATTCATTAG	8
sep	AATCATAACCAACCGAATCA TCATAATGGAAGTGCTATAA	8
sea	AGATCATTCGTGGTATAACG TTAACCGAAGGTTCTGTAGA	8
mecA	TCCAGATTACAACTTCACCAGG CCACTTCATATCTTGTAACG	14

Table 2) Different IEC phage types and their incidencefound among the MRSA isolates

Phage type	Genes	Number (%)
А	sea, sak, chp, scn	7 (4.8)
В	sak, chp, scn	13 (9)
С	chp, scn	19 (13.1)
D	sea, sak, scn	18 (12.4)
Е	sak, scn	40 (27.6)
F	sep , sak, chp, scn	2 (1.4)
G	sep , sak, scn	1 (0.7)
Н	scn	10 (6.9)
None	sep or sak or chp or sea or none	35 (24.1)
Total		145 (100)

Ethics Statement: All the study stages complied with the Declaration of Helsinki. Also, oral consents were obtained from all the participants for the participation in this study.

Statistical analysis: Fisher's test and Chisquare test were used for the comparison of data. For all statistical analyses, SPSS software (v.22.0) was used. P-values less than 0.05 were considered as statistically significant.

Findings

Clinical features of *S. aureus* **isolates:** Out of 145 MRSA isolates, 67 (46.2%) isolates belonged to the female participants, while 78 (53.8%) strains were isolated from male participants (P> 0.05). The strains were mostly isolated from wound (46.2%), followed by blood (15.9%), respiratory infection (13.1%), urine (11%), catheter (7.6%), and abscess (6.2%) samples.

Antimicrobial susceptibility profiles of *S. aureus* isolates: All the isolates

Table 3) Antibiotic resistance patterns of MRSA isolates

Sensitive

N (%)

4 (2.75)

Antimicrobial agent

Р

were assessed for susceptibility to 10 antibiotics. The antimicrobial resistance patterns of the strains are listed in Table 3. All the investigated isolates were cefoxitinresistant and *mecA* gene positive. Among the isolates, the highest resistance rate was recorded towards penicillin (97.25%), followed by methicillin (95.8%), ceftazidime (81.4%), and erythromycin (71.8%). None of the isolates were resistant to vancomycin. On the other hand, MDR (\geq 3 classes) was recorded in 95.5% of the isolates.

IEC genes and phage typing of the isolates: Among the IEC genes, *scn* gene was the most prevalent (110; 75.8%), followed by *sak* (87; 60%), *chp* (42; 28.9%), *sea* (27; 18.6%), and *sep* (4; 2.75%) genes. Eight types (A-H) were previously defined for different combinations of IEC genes, harbored by β -hemolysinconverting bacteriophage. The predominant IEC scheme was Type E. Overall, 35 isolates showed no IEC type and contained none or only one gene. The details of the detected types are presented in Table 3.

Resistant

141 (97.25)

N (%)

	. ,		. ,
V	135 (93.1)	10 (6.9)	0 (0)
СР	51 (35.2)	6 (4.1)	88 (60.7)
Е	32 (22)	9 (6.2)	104 (71.8)
CC	53 (36.5)	3 (2.1)	89 (61.4)
GM	61 (42)	3 (2)	81 (56)
CAZ	11 (7.6)	16 (11)	118 (81.4)
IPM	59 (40.7)	4 (2.75)	82 (56.55)
Ме	4 (2.8)	2 (1.4)	139 (95.8)
Fox	0 (0)	0 (0)	145 (100)

Intermediate

N (%)

0(0)

P: penicillin; V: vancomycin; CP: ciprofloxacin; E: erythromycin; CC: clindamycin; GM: gentamicin; CAZ: ceftazidime; IMP: imipenem; Me: methicillin; and Fox: cefoxitin

Discussion

S. aureus is an important human and livestock pathogen, which can cause a broad spectrum of diseases ^[7]. The rapid growing resistance of this bacterium to antibiotics, such as methicillin and other β -lactam antibiotics, imposes challenges to the infection treatment. The prevalence of MRSA is higher in Iran than in neighboring countries in the Middle East (except for Iraq) ^[15, 17]. Evidence suggests that the frequency of some virulence genes in MRSA isolates is higher than in methicillin-susceptible S. aureus (MSSA) strains; this can be the reason for great diversity and variability in these isolates ^[2, 18, 19]. In MRSA isolates, the IEC genes carried by β -hemolysinconverting bacteriophages, facilitate nasal colonization and adaptation of S. aureus isolates to their human host [9, 20, 21]; in addition, propagation and intracellular survival of bacteria increase in the nose ^[9]. However, different IEC types are usually carried by bacteriophages. It should be noted that bacteria usually tend to lose their bacteriophages, which is because of variable frequency of IEC genes among the S. aureus isolates ^[22].

The antimicrobial susceptibility test of MRSA isolates indicated that penicillin, methicillin, ceftazidime, erythromycin, clindamycin, and ciprofloxacin are not effective empirical options in the treatment of S. aureus infections. Also, resistance to these antimicrobial drugs has been reported in other studies from Iran; therefore, empirical treatment of MRSA-infected patients with these antibiotics may not be effective ^[15, 23, 24]. Nevertheless, vancomycin can be an effective antibiotic against MRSA, as all the tested isolates were susceptible to this antimicrobial drug. This finding is in agreement with the finding of other studies conducted in Iran^[15, 25]. In the present study, more than 95% of the S. aureus isolates

were MDR, indicating the widespread antimicrobial resistance in *S. aureus* isolates and the need for continuous monitoring the antibacterial resistance development in MRSA isolates.

In the present study, 75.9% of MRSA isolates contained an IEC-carrying bacteriophage, which was slightly lower than that reported in other studies ^[8, 9]. However, lower prevalence rate was reported in another study^[1], which is presumably related to the animal origin of these isolates. In another study conducted in Iran, the prevalence of IEC-carrying bacteriophages among the clinical S. aureus isolates was lower than that reported in the present study ^[15]. In addition, several differences were observed in the prevalence of phage types in the clinical MRSA isolates, compared to those reported in some previous studies. The predominant IEC variant was Type E, while in other studies, human infecting isolates mostly belonged to Type B [3, 8, 9, ^{15]}. These findings suggest the presence of variations in IEC types, found in the isolates of different geographical regions; this stems from the fact that IEC is a dynamic DNA element, which can successfully spread among the S. aureus isolates, resulting in the adaptation and counteraction to the human host [8].

Although in this study, prophage integration was found in approximately 76% of the human clinical isolates, the prevalence of individual genes was varied, which is due to the fact that all IEC genes are not transferred by a given bacteriophage. Similar to a study by van Wamel et al. (2006), the most frequent genes were scn, sak, and chp genes, respectively [8]. Nonetheless, the prevalence of *chp* gene was varied considerably between the different studies, indicating the reduced prevalence of phage Type B and increased prevalence of phage Type E.

Conclusion

In conclusion, based on the reported antimicrobial susceptibility testing results, vancomycin can be considered as an effective empirical option for the treatment of MRSA infections. The IEC-carrying bacteriophages are highly prevalent among the MRSA strains and varied in different geographical regions. This reveals the successful and continuous spread of IEC through the clinical *S. aureus* population. Therefore, IEC genes can act as the potential targets for the vaccination or treatment of *S. aureus* infections.

Acknowledgements: This study is related to the MS thesis by Simin Ariyarad, approved by the Shahid Chamran University of Ahvaz. Authors are very thankful to Shahid Chamran University of Ahvaz for providing facilities to accomplish the present research project at the specified time.

Ethical Permissions: The study was approved by the research committee of Shahid Chamran University of Ahvaz. **Conflicts of interest:** The authors declare that they have no competing interest.

Authors' contribution: SER and SA participated in search for subjects and data, also performance of the research. SER conducted the research. SER and HM analyzed and interpreted the data. All authors read, revised, and approved the final manuscript.

Funding: This study was supported by a grant from Shahid Chamran University of Ahvaz (No: 95/3/02/31400), Iran.

References

1. Baptistão LG, Silva NC, Bonsaglia EC, Rossi BF, Castilho IG, Fernandes Junior A, et al. Presence of immune evasion cluster and molecular typing of methicillin-susceptible Staphylococcus aureus isolated from food handlers. J Food Prot. 2016; 79(4):682-6. 2. Liang BS, Huang YM, Chen YS, Dong H, Mai JL, Xie YQ, et al. Antimicrobial resistance and prevalence of CvfB, SEK, and SEQ genes among Staphylococcus aureus isolates from pediatric patients with bloodstream infections. Exp Ther Med. 2017; 14(5):5143-8.

3. Hau SJ, Sun J, Davies PR, Frana TS, Nicholson TL. Comparative prevalence of immune evasion complex genes associated with β -hemolysin converting bacteriophages in MRSA ST5 isolates from swine, swine facilities, humans with swine contact, and humans with no swine contact. PLoS One. 2015; 10(11):e0142832.

4. McCarthy AJ, Witney AA, Lindsay JA. Staphylococcus aureus temperate bacteriophage: Carriage and horizontal gene transfer is lineage associated. Front Cell Infect Microbiol. 2012; 2:6.

5. Foster TJ. Immune evasion by staphylococci. Nat Rev Microbiol. 2005; 3(12):948-58.

6. Yu F, Li T, Huang X, Xie J, Xu Y, Tu J, et al. Virulence gene profiling and molecular characterization of hospital-acquired Staphylococcus aureus isolates associated with bloodstream infection. Diagn Microbiol Infect Dis. 2012; 74(4):363-8.

7. Jongerius I, von Köckritz-Blickwede M, Horsburgh MJ, Ruyken M, Nizet V, Rooijakkers SH. Staphylococcus aureus virulence is enhanced by secreted factors that block innate immune defenses. J Innate Immun. 2012; 4(3):301-11.

8. van Wamel WJ, Rooijakkers SH, Ruyken M, van Kessel KP, van Strijp JA. The innate immune modulators staphylococcal complement inhibitor and chemotaxis inhibitory protein of Staphylococcus aureus are located on beta-hemolysin-converting bacteriophages. J Bacteriol. 2006; 188(4):1310-5.

9. Verkaik NJ, Benard M, Boelens HA, de Vogel CP, Nouwen JL, Verbrugh HA, et al. Immune evasion cluster-positive bacteriophages are highly prevalent among

DOR: 20.1001.1.25884107.2019.5.1.2.3

human Staphylococcus aureus strains, but they are not essential in the first stages of nasal colonization. Clin Microbiol Infect. 2011; 17(3):343-8.

10. Jongerius I, Köhl J, Pandey MK, Ruyken M, van Kessel KP, van Strijp JA, et al. Staphylococcal complement evasion by various convertase-blocking molecules. J Exp Med. 2007; 204(10):2461-71.

11. Kraushaar B, Hammerl JA, Kienöl M, Heinig ML, Sperling N, Dinh Thanh M, et al. Acquisition of virulence factors in livestockassociated MRSA: Lysogenic conversion of CC398 strains by virulence gene-containing phages. Sci Rep. 2017; 7(1):2004.

12. de Jong NWM, Vrieling M, Garcia BL, Koop G, Brettmann M, Aerts PC, et al. Identification of a staphylococcal complement inhibitor with broad host specificity in equid Staphylococcus aureus strains. J Biol Chem. 2018; 293(12):4468-4477.

13.Wayne PA. CLSI, Performance standards for antimicrobial susceptibility testing; Twenty-second informational supplement. CLSI document M100-S22. Clinical and Laboratory Standards Institute, 2017.

14. Emaneini M, Bigverdi R, Kalantar D, Soroush S, Jabalameli F, Noorazar Khoshgnab B, et al. Distribution of genes encoding tetracycline resistance and aminoglycoside modifying enzymes in Staphylococcus aureus strains isolated from a burn center. Ann Burns Fire Disasters. 2013;26(2):76-80. 15. Ahmadrajabi R, Layegh-Khavidaki S, Kalantar-Neyestanaki D, Fasihi Y. Molecular analysis of immune evasion cluster (IEC) genes and intercellular adhesion gene cluster (ICA) among methicillin-resistant and methicillin-sensitive isolates of Staphylococcus aureus. J Prev Med Hyg. 2017;58(4): E308-14.

16. Price LB, Stegger M, Hasman H, Aziz M, Larsen J, Andersen PS, et al. Staphylococcus aureus CC398: host adaptation and emergence of methicillin resistance in livestock. MBio. 2012; 3(1): e00305-11. 17. Stefani S, Chung DR, Lindsay JA, Friedrich AW, Kearns AM, Westh H, et al. Meticillinresistant Staphylococcus aureus (MRSA): global epidemiology and harmonisation of typing methods. Int J Antimicrob Agents. 2012; 39(4):273-82.

18. Hudson LO, Murphy CR, Spratt BG, Enright MC, Elkins K, Nguyen C, et al. Diversity of methicillin-resistant Staphylococcus aureus (MRSA) strains isolated from inpatients of 30 hospitals in Orange County, California. PLoS One. 2013; 8(4):e62117.

19. Dinges MM, Orwin PM, Schlievert PM. Exotoxins of Staphylococcus aureus. Clin Microbiol Rev. 2000; 13(1): 16-34.

20.You Y, Song L, Nonyane BAS, Price LB, Silbergeld EK. Genomic differences between nasal Staphylococcus aureus from hog slaughterhouse workers and their communities. PLoS One. 2018; 13(3):e0193820.

21.Zadoks RN, van Leeuwen WB, Kreft D, Fox LK, Barkema HW, Schukken YH, et al. Comparison of Staphylococcus aureus isolates from bovine and human skin, milking equipment, and bovine milk by phage typing, pulsed-field gel electrophoresis, and binary typing. J Clin Microbiol. 2002; 40(11):3894–902.

22.Goerke C, Wirtz C, Fluckiger U, Wolz C. Extensive phage dynamics in Staphylococcus aureus contributes to adaptation to the human host during infection. Mol Microbiol. 2006; 61(6):1673-85.

23.Sadeghi J, Mansouri S. Molecular characterization and antibiotic resistance of clinical isolates of methicillin-resistant Staphylococcus aureus obtained from Southeast of Iran (Kerman). APMIS 2014; 122(5):405-11.

24.Ohadian Moghadam S, Pourmand MR, Mahmoudi M, Sadighian H. Molecular characterization of methicillin-resistant Staphylococcus aureus: Characterization of

DOR: 20.1001.1.25884107.2019.5.1.2.3

major clones and emergence of epidemic clones of sequence type (ST) 36 and ST 121 in Tehran, Iran. FEMS Microbiol Lett. 2015; 362(8):fnv043.

25. Saderi H, Emadi B, Owlia P. Phenotypic and

genotypic study of macrolide, lincosamide, and streptogramin B (MLSB) resistance in clinical isolates of Staphylococcus aureus in Tehran, Iran. Med Sci Monit. 2011; 17(2):BR48-53.