

of which 5 isolates were from EKSUTH samples, and 2 isolates were from SSHIE samples. ARG2b had 2 resistant isolates, which were only from EKSUTH samples with a total occurrence of 45%.

Antibiotics inhibition: Figure 2 shows the similarity between the antibiotics reactions with respect to the resistance behavior of 20 *S. aureus* isolates. The clusters were made up of 3 different antibiotic groups, namely Group 1 (Grp1), Group 2 (Grp2), and Group 3 (Grp3) based on the reaction of the 8 antibiotics against the 20 *S. aureus* isolates. Antibiotics of Group 1 included OFL, GENT, and ERY. Antibiotics of Group 2 included CRX and CTR; and antibiotics of Group 3 included AUG, CAZ, and CXC. Antibiotics OFL and GENT which were both of Group 1 behaved in the same way in their reaction

so that a high level of isolates susceptibility was observed to these antibiotics; however, isolates expressed a similar resistance behavior in their reaction to CAZ and CXC antibiotics both of Group 3. Antibiotics 1 and 2 of Grp1 and Antibiotics 3 and 4 of Grp3 were identical in their reaction on the 20 *S. aureus* isolates with 25% similarity.

Plants' extracts inhibition: Figure 3 presenting cluster E shows how the leaves extracts of *T. catappa*, *M. indica*, and *A. wikesiena* inhibited 20 isolated *S. aureus* strains. Two major groups of ERG1 and ERG2 were formed by 55% coefficient of similarity. From the first group, 2 subgroups were emerged, including ERG1a and ERG1b. ERG1a consisted of 12 isolates including 1-MSWB5U, 2-FSWB4U, 3-FSWB4iiU, 4-MMWB1iU, 17-FMWB3i, 18-MSWB6iiWS,




Figure 2) Similarity between the antibiotics reactions with respect to the resistance behavior of 20 *S. aureus* isolates

the findings of Onanuga et al. (2012) [11]. Modern science and technology have achieved success in new drug discovery; nevertheless, pathogens with high frequency develop means to resist against some synthesized drugs used for their treatment. Extensive use of broad-spectrum antibiotics results in the widespread prevalence of nosocomial infections caused by multidrug resistant pathogens (Chikere et al. (2008)[26]. Therefore, the high prevalence of *S. aureus* strains in this study and their resistance pattern to most of the antibiotics used make it necessary to pay needed attention in order to aware the world about the adverse effects of multidrug resistant microorganisms. This aim could be achieved by finding new measures to combat them. However, some ineffective synthesized drugs are of more recognition than those faced with in traditional folklore. The prescription and use of these ineffective drugs may favor the emergence of hospital and community acquired resistant strains of pathogens. Studied have revealed that resistance has been almost observed to all antibiotic groups [27]. In addition, adverse side effects of some antibiotics on the host have been reported in many studies, including hypersensitivity, depletion of beneficial gut and mucosal microorganisms, immunosuppression, and allergic reactions [28], those which have not been reported for plant extracts. The number of multi-drug resistant microbial strains and the emergence of strains with reduced susceptibility to antibiotics are continuously increasing. This increase has been attributed to the indiscriminate use of broad-spectrum antibiotics, immunosuppressive agent, intravenous catheters, organ transplantation, and the epidemics of HIV infection [29, 30]. Among the 20 isolated *S. aureus* species, some were found to be resistant only to *A. wikesiena*, among the employed plant extracts. This suggests that the isolates

need to be exposed to a wide range of plant extracts instead of only 3, compared to the 8 antibiotics employed. Doing so would produce a better result in comparison with the existing antibiotics and would be a solution for discovering alternatives to many antibiotics that are ineffective on *S. aureus* origin infections.

In comparison, the inhibitions recorded with standard reference antibiotics were comparable to those observed with various leaves extracts; although some of the antibiotics such as OFL, GEN, ERY, CRX, GEN, and ERY exhibited higher inhibitory effect on the tested isolates, compared to *A. wikesiena* leaf extract.

Among the antibiotics used, OFL inhibited most the *S. aureus* isolates growth. This apparently high level of susceptibility to OFL suggests that it could be considered as a drug of choice for treating infections caused by *S. aureus* in the study area, especially at the present time that *S. aureus* strains are resistant to many commonly used antibiotics. Similar result about the high susceptibility of *S. aureus* strains to OFL was reported by Uwaezuoke and Aririatum (2004) [31].

The isolated *S. aureus* strains were resistant to CXC, CAZ, AUG, and ERY. The high level of resistance observed could be due to early exposure of these drugs to the isolates which may have enhanced resistance. This suggests that there is a high level of antibiotics abuse in the studied areas, which could be due to self-medication, inadequate dosage, and failure to comply with treatment, and availability of antibiotics to consumers across the counters with or without prescription, as reported by Odugbemi (1981) [32].

Identification of antibiotic resistant isolates using molecular techniques such as 16S rRNA, Random Amplification of Polymorphic DNA (RAPD), and plasmid profile causing infectious processes is usually essential for effective antimicrobial and supportive

therapy^[33]. Initial treatment may be empirical based on the microbiological epidemiology of the infection and the patient's symptoms^[34]. From the epidemiological point of view, in this study, it was observed that all the strains isolated from patients receiving treatment from EKSUTH exhibited a similar behavior in their resistance to the antibiotics used. There was a relationship between the ERG and ARG *S. aureus* isolates population structure across the three studied locations in Ekiti State.

The use of molecular methods (RAPD- PCR) confirmed the obtained results and that the isolates responding similarly to certain antibiotics and plant extracts shared similar genotypes and should therefore be grouped in the same clusters. By virtue of this analysis, it was possible to group isolates with similar genotypes and similar resistance behavior to the antibiotics and plants extracts in the same clusters.

Conclusions

Based on the obtained results in this study, it could be deduced that *S. aureus* infection is high and common among the patients. Infection rate was higher in the age groups below 50 years, and the risk factors were mainly poor hygiene and socio economic conditions. The differences observed in the colonization rates between the two sexes indicated that sex could be considered as a risk factor for *S. aureus* infection among patients. This was the first study reporting the epidemiology of *S. aureus* strains isolated from patients attending to hospital treatments in Ado Ekiti, Nigeria. This study provided information about the *S. aureus* infection rate in different age groups of patients receiving hospital treatments and also about the employed plant extracts for effective treatment of *S. aureus* infections, in addition to the existing traditional medication.

Acknowledgement: The authors are very grateful to all the health workers in Ekiti State University Teaching Hospital, State Specialist Hospital Ekere-Ekiti, and General Hospital Iyin-Ekiti, assisting in the samples collection. The authors would like to appreciate Prof. Onasanya Amos for his assistance in samples analysis.

Ethical Permissions: The protocol was confirmed in ethical committee, and the authors of this manuscript observed ethical issues (EKSUTH/A67/2016/10/021)

Conflict of interests: The authors declare that there is no conflict of interest.

Authors' contribution: MOO and FCA conceived the research idea, designed, and wrote the first draft of the manuscript. FCA managed the literature search. All authors read and approved the final manuscript.

Fundings: None.

References

1. Weber DJ, Rutala WA. Self-disinfecting surfaces: Review of current methodologies and future prospects. *Am J Infect Control*. 2013; 41(5):S31–5.
2. Mathur P. Hand hygiene: Back to the basics of infection control. *Indian J Med Res*. 2011; 134(5):611–20.
3. Shinagawa N, Taniguchi M, Hirata K, Furuhashi T, Fukuhara K, Mizuguchi T. Bacteria isolated from surgical infections and its susceptibilities to antimicrobial agents-special references to bacteria isolated between April 2010 and March 2011. *Japan J Antibiot*. 2014; 67(5):293–334.
4. Harris L, Foster S, Richards S, An Hannan Abdul, Asghar Samra, Naeem Tahir Ullah, Muhammad Ikram. Medicinal effects of *Mangifera indica*. *Pak Jour Pharm Sci*. 2002;26:4-715.
5. Elkins C, Thomas CE, Selfert HS, Sparling PF. Species-specific uptake of DNA by Gonococci is mediated by

- a 10-base-pair sequence. *J Bacteriol.* 1999; 173(12):3911-3.
6. Onasanya A, Mignouna HD, Thottappilly G. Genetic fingerprinting and phylogenetic diversity of isolates of *Staphylococcus aureus* from Nigeria. *Afr J Biotechnol.* 2003; 2(8):246-50.
 7. Rohlf FJ. NTSYS-pc: numerical taxonomy and multivariable analysis system. State university of New York: Stony brooks; 1989.
 8. Clinical and Laboratory Standards Institute (CLSI). M02-A11: Performance standards for antimicrobial disc susceptibility tests; Approved standard. 7th Edition. Wayne: Clinical and Laboratory Standards Institute; 2012.
 9. Orenstein R, Wong ES. Urinary tract infections in adults. *Am Fam Physician.* 1999; 59(5):1225-34.
 10. Khan SW, Ahmed A. Uropathogens and their susceptibility pattern: A retrospective analysis. *J Pak Med Assoc.* 2001; 51(2):98-9.
 11. Onanuga A, Awhowho GO. Antimicrobial resistance of *Staphylococcus aureus* strains from patients with urinary tract infections in Yenagoa, Nigeria. *J Pharm Bioallied Sci.* 2012; 4(3):226-30.
 12. Nwanze PI, Nwaru LM, Oranusi S, Dimkpa U, Okwu MU, Babatunde BB. Urinary tract infection in Okada village: Prevalence and antimicrobial susceptibility pattern. *Sci Res Essay.* 2007; 2(4):112-6.
 13. Akortha EE, Ibadin OK. Incidence and antibiotic susceptibility pattern of *Staphylococcus aureus* amongst patients with urinary tract infection (UTIS) in UBTH Benin City, Nigeria. *Afr J Biotechnol.* 2008; 7(11):1637-40
 14. Aboderin OA, Abdu A, Odetoyin BW, Lamikanra A. Antimicrobial resistance in *Escherichia coli* strains from urinary tract infections. *J Natl Med Assoc.* 2009; 101(12):1268-73.
 15. Akerele J, Akonkhai I, Isah A. Urinary pathogens and antimicrobial susceptibility: A retrospective study of private diagnostic laboratories in Benin City, Nigeria. *J Med Lab Sci.* 2000; 9:47-52
 16. Okonko IO, Donbraye-Emmanuel OB, Ijandipe LA, Ogun AA, Adedeji AO, Udeze AO. Antibiotics sensitivity and resistance patterns of uropathogens to nitrofurantoin and nalidixic acid in pregnant women with urinary tract infections in Ibadan, Nigeria. *Middle-East J Sci Res.* 2009; 4(2):105-9.
 17. Manikandan S, Ganesapandian S, Singh M, Kumaraguru AK. Antimicrobial susceptibility pattern of urinary tract infection causing human pathogenic bacteria. *Asian J Med Sci.* 2011; 3(2):56-60.
 18. Deyno S, Toma A, Worku M, Bekele M. Antimicrobial resistance profile of *Staphylococcus aureus* isolates isolated from ear discharges of patients at University of Hawassa comprehensive specialized hospital. *BMC Pharmacol Toxicol.* 2017; 18(1):35.
 19. Almagadam BS, Ahmed HM, Osman M.B, Omer SM. Frequency of MRSA isolates in mobile phones, ears, and hands of healthcare workers. (2018). *J Antimicrob Agents.* 2018; 4(1): 161.
 20. Kuehnert MJ, Deanna K, Hill HA. Prevalence of *Staphylococcus aureus* nasal colonization in the United States. *J Infect Dis.* 2006; 193(2):172-9.
 21. Plat J, Rai SK. Occurrence of *Staphylococcus aureus* in hospital environment and staffs in teaching hospital in Katmandu, Nepal. *J Nepal Assoc Med Lab Sci.* 2007; 8(1):72-3.
 22. Chigbu CO, Ezeronye OU. Antibiotic resistant *Staphylococcus aureus* in Abia State of Nigeria. *Afr J Biotechnol.* 2003; 2(10): 374-8.

23. Adesina SK, Idowu O, Ogundaini AO, Oladimeji H, Olugbade TA, Onawunmi GO. Antimicrobial constituents of the leaves of *Acalypha wilkesiana* and *Acalypha hispida*. *Phytother Res*. 2000; 14(5): 371-4.
24. Mulu W, Kibru G, Beyene G, Dامتie M. Postoperative nosocomial infections and antimicrobial resistance pattern of bacteria isolates among patients admitted at Felege Hiwot referral hospital, Bahirdar, Ethiopia. *Ethiop J Health Sci*. 2012; 22(1):7-18.
25. Muluye D, Wondimeneh Y, Ferede G, Nega T, Adane K, Biadgo B, et al. Bacterial isolates and their antibiotic susceptibility patterns among patients with pus and/or wound discharge at Gondar university hospital. *BMC Res Notes*. 2014; 7(1):619.
26. Chikere CB, Chikere BO, Omoni VT. Antibiogram of clinical isolates from a hospital in Nigeria. *Afr J Biotechnol*. 2008; 7(24): 4359-63.
27. Eloff JN. On expressing the antibacterial activity of plant extracts-a small first step in applying scientific knowledge to rural primary health care. *South Afr J Sci*. 2000; 96(3): 16-118.
28. Al-Jabri AA. Honey, milk, and antibiotics. *Afr J Biotechnol*. 2005; 4(13):1580-7.
29. Dean DA, Burchard KW. Fungal infection in surgical patients. *Am J Surg*. 1996; 171(3):374-82.
30. Gonzalez CE, Venzon D, Lee S, Mueller BU, Pizzo PA, Walsh TJ. Risk factors for fungemia in children infected with human immunodeficiency virus: A case-control study. *Clin Infect Dis*. 1996; 23(3): 515-21.
31. Uwaezuoke JC, Aririatu LE. A survey of antibiotic resistant *Staphylococcus aureus* strains from clinical sources in Owerri. *J Appl Sci Environ Managt*. 2004; 8(1):67-8.
32. Odugbemi T. The use and abuse of antibiotics. *Nig Med Pract*. 1981; 1(1):4-8.
33. Adesida SA, Abioye OA, Bamiro BS. Associated risk factors and pulsed field gel electrophoresis of nasal isolates of *Staphylococcus aureus* from medical students in a tertiary hospital in Lagos, Nigeria. *Braz J Infect Dis*. 2007; 11(1):63-9.
34. Al-Khulaifi M, Amin Aref M, Al Salamah A. Phage typing, PCR amplification for *mecA* gene, and antibiotic resistance patterns as epidemiologic markers in nosocomial outbreaks of methicillin resistant *Staphylococcus aureus*, Saudi J Biol Sci. 2009; 16 (1):37-49.