

# Synergistic Effect of *Artemisia scoparia* Extract and TiO<sub>2</sub> Nanoparticles on Antibiotic Resistance of *Klebsiella pneumoniae*

## ARTICLE INFO

Article Type  
Original Research

### Authors

Saharnazsadat Serajian, MSc<sup>1</sup>  
Mina Ramezani, PhD<sup>2\*</sup>  
Sahar Honarmand Jahromy, PhD<sup>3</sup>

### How to cite this article

Serajian S.S., Ramezani M., Honarmand Jahromy S. Synergistic Effect of *Artemisia scoparia* Extract and TiO<sub>2</sub> Nanoparticles on Antibiotic Resistance of *Klebsiella pneumoniae*. Infection Epidemiology and Microbiology. 2021;7(2): 101-108

<sup>1</sup> Department of Biology, Central Tehran Branch, Islamic Azad University, Tehran, Iran

<sup>2</sup> Department of Biology, Central Tehran Branch, Islamic Azad University, Tehran, Iran

<sup>3</sup> Department of Microbiology, Varamin Branch, Islamic Azad University, Varamin, Iran

### \* Correspondence

Address: Department of Biology, Central Tehran Branch, Islamic Azad University, Imam Hasan Bld, Ponak Square, Tehran, Iran.  
m.ramezani@iauctb.ac.ir

### Article History

Received: March 05 2021  
Accepted: April 25, 2021  
Published: May 20, 2021

## ABSTRACT

**Backgrounds:** This study aimed to assess antibacterial properties of *Artemisia scoparia*, Titanium dioxide nanoparticles, and their synergistic effect on clinical isolates of *Klebsiella pneumoniae*.

**Materials & Methods:** In this experimental study, 30 isolates of *K. pneumoniae* were collected from patients' sputum in the microbiology lab of Masih Daneshvari hospital during 3 months. Then biochemical tests were performed for strain confirming. Moreover, genomic DNA was extracted from all the isolates, and *hly* gene was detected in the isolates via PCR method. The susceptibility of the isolates to 10 antibiotics was evaluated by the disk diffusion method. Then minimum inhibitory concentration (MIC) of all components (*Artemisia* extract, TiO<sub>2</sub>, and their combination) was assessed using the microdilution method against the isolates.

**Findings:** The results indicated that simultaneous use of hydro-alcoholic extract of *A. scoparia* and titanium dioxide nanoparticles exhibited a significant synergistic antibacterial effect on 25 clinical isolates in comparison with the use of extract or nanoparticles alone.

**Conclusion:** It seems that simultaneous use of *Artemisia* herbal extracts and nanoparticles is beneficial in increasing their antibacterial effect and may decrease antibiotics consumption.

**Keywords:** *Klebsiella pneumoniae*, Titanium dioxide, Nanoparticles, *Artemisia scoparia*.

## CITATION LINKS

[1] Adams-Sapper S, Nolen S, Donzelli GF, Lal M, Chen K, Justo da Silva LH, et al. Rapid induction of ... [2] Bengoechea JA, Sa Pessoa J. *Klebsiella pneumoniae* infection ... [3] Karabinis A, Paramythiotou E, Mylona-Petropoulou D, Kalogeromitros A, Katsarelis N, Kontopidou F, et al. Colistin for *Klebsiella pneumoniae*-associated... [4] Soltan Dalal MM, Miremadi SA, Sharifi Yazdi MK, Rastegar Lari AA, Rajabi Z, Avadis YS. Antimicrobial ... [5] Behzadian Nejad Q, Abdollahi A, Najar Peerayeh SH, Forouhesh ... [6] Wyres KL, Lam MM, Holt KE. Population genomics of *Klebsiella* ... [7] Shakeel M, Jabeen F, Shabbir S, Asghar MS, Khan MS, Chaudhry AS. Toxicity of ... [8] Shi H, Magaye R, Castranova V, Zhao J. Titanium dioxide ... [9] Chellappa M, Anjaneyulu U, Manivasagam G, Vijayalakshmi U. Preparation and ... [10] Garcia-Contreras R, Scougall-Vilchis RJ, Contreras-Bulnes R, Ando Y, Kanda Y, Hibino Y, et al. Effects of ... [11] Khan ST, Ahmad J, Ahamed M, Jousset A. Sub-lethal doses of widespread ... [12] Nam SY, Han NR, Rah SY, Seo Y, Kim HM, Jeong HJ. Anti-inflammatory effects of *Artemisia* ... [13] Ramezani M, Fazli-Bazzaz BS, Saghafi-Khadem F, Dabaghian A. Antimicrobial activity of ... [14] Sajid M, Rashid Khan MR, Shah NA, Waris TS, Younis T, Ullah S, et al. Evaluation of *Artemisia scoparia* for ... [15] Cha JD, Jeong MR, Jeong SI, Moon SE, Kim JY, Kil BS, et al. Chemical composition and antimicrobial activity of the essential oils of *Artemisia scoparia* and *A. capillaris*. *Planta* ... [16] Singh HP, Mittal S, Kaur S, Batish DR, Kohli RK. Chemical composition and antioxidant activity of ... [17] Candan ED, Aksöz N. *Klebsiella pneumoniae*: Characteristics of carbapenem resistance and ... [18] Singh J. Maceration, percolation, and infusion techniques for the ... [19] Martin RM, Bachman MA. Colonization, infection, and the accessory genome of *Klebsiella* ... [20] Ayati R, Dodi M, Karami M. Evaluation the effect of TiO<sub>2</sub> nanoparticles on MDR *Klebsiella pneumoniae* ... [21] Verdier T, Coutand M, Bertron A, Roques C. Antibacterial activity of TiO<sub>2</sub> photocatalyst alone or ... [22] Tambekar DH, Dahikar SB. Antibacterial activity of some Indian Ayurvedic preparations against ... [23] Winnett V, Sirdaarta J, White A, Clarke FM, Cock IE. Inhibition of *Klebsiella pneumoniae* growth by selected Australian plants: Natural approaches for the prevention and management of ankylosing spondylitis. *Inflammopharmacology*. 2017;25(2):223-35. [24] Gavanji S, Mohammadi E, Larki B, Bakhtari A. Antimicrobial and cytotoxic... [25] Haroun MF, Al-Kayali RS. Synergistic effect of *Thymbra spicata* L. extracts... [26] Hacıoğlu M, Dosler S, Birteksoz Tan AS, Otuk G. Antimicrobial activities of widely consumed herbal teas, alone or in combination with...

## Introduction

*Klebsiella* is a Gram-negative bacterium of the large family *Enterobacteriaceae*, which is very common in different communities. This bacterium exists as a saprophyte microorganism in human nasopharynx and intestine and is considered as an opportunistic bacterium one of the main causes of pneumonia, nosocomial infections, and urinary tract infection (UTI). In recent years, *klebsiella* has been reported to cause many infections, and the importance of this group of organisms as a cause of serious infections in hospitalized patients has been accepted [1-2]. The ability of this organism to cause diseases due to weakened host defenses as a result of complex and lengthy surgeries as well as the use of various antibiotics has been well documented [3]. The rate of *Klebsiella* colonization is 2 to 4 times higher in hospitalized patients, especially in patients taking broad-spectrum antibiotics [4]. This strain is naturally resistant to penicillin and could acquire resistance to several antimicrobials [5-6]. Today, due to the drug resistance of *K. pneumoniae* to several antibiotics (Multiple Drug resistance) from different classes, including penicillin and carbapenem, there is a need to use new and combination therapies against this strain. In the last decade, the use of nanomaterials has become increasingly widespread so that they have entered all aspects of life. Meanwhile, the use of nanomaterials in medical processes has also increased [7-8]. Titanium dioxide (TiO<sub>2</sub>) nanoparticles are an important product in nanotechnology, which are used as a pharmaceutical compound and an attractive candidate to deliver many small drug molecules or large biomolecules such as DNA, RNA, and proteins. TiO<sub>2</sub> nanoparticles are heat resistant and could emit O<sup>2-</sup> and OH<sup>-</sup> species due to light radiation. These compounds could affect membrane lipids and cause their degradation [9-11].

One of the popular treatment approaches for microbial infections is the use of plant extracts. *Artemisia* belongs to the *Asteraceae* (*Compositae*) family. This plant includes 200 to 500 species or subspecies and 5 subgenera. Iran has 34 species of *Artemisia* which is one of the most important plant species after *Astragalus* in terms of distribution and density. Therefore, *Artemisia* community has a 60% coverage [12]. This genus includes several types of medicinal plants. Different species of this genus are used not only in the pharmaceutical industry but also in the food and cosmetics industries. *Artemisia* is rich in substances that have various effects including anti-inflammatory, anti-tumor, anti-ulcer, diuretic, antioxidant, anti-malarial, indigestion relieving, anti-cell proliferation, and antimicrobial properties [13-14]. *A. scoparia* contains essential oils that have been reported to have antimicrobial and antioxidant effects [15-16].

**Objectives:** Given the dangerous diseases caused by *K. pneumoniae* and the prevalence of MDR strains, it is important to treat *K. pneumoniae* infections by new compounds. The aim of this study was to evaluate the synergistic antibacterial effect of *A. scoparia* extract and TiO<sub>2</sub> nanoparticles on clinical isolates of *K. pneumoniae*.

## Materials and Methods

In this experimental study, clinical samples were collected from patients with suspected respiratory infection, referring to Masih Daneshvari hospital during 3 months in autumn and winter of 2020. The best sample was morning sputum, which was collected through a deep cough. All ethical considerations were taken into account in the research, and informed written consent was obtained from the patients. Samples were cultured on Blood Agar, EMB Agar, and McKankey (Merck, Germany) plates. Subculturing and purification of colonies with similar properties

to *Klebsiella* were performed after 24-48 hours of incubation at 37 °C. Gram staining, differential culture medium, and biochemical tests including indole, Simmon citrate motion, triple-sugar iron agar (TSI), methyl red (MR), and urease (Merck, Germany) were used to identify *K. pneumoniae* isolates.

**Molecular identification of *K. pneumoniae*:**

To confirm the phenotypic tests, PCR reaction was performed to amplify the 77 bp region of the *hly* gene encoding hemolysin which is specific for this strain. To extract DNA, a DNA purification kit (Biospin Bacteria Genomic DNA Extraction Kit, Bioer) was used. Materials used for PCR reaction were as follows: DNA 1 µL (50 ng/µL), Master Mix 12.5 µL, and Primer Mix 2 µL, and the final volume was increased to 25 µL by deionized water. Specific forward and reverse primers (PF: GATGAAACGACCTGATTGCATTC and PR: CCGGGCTGTCGGGATAAG) were used [17]. Temperature cycles included: primary denaturation at 95 °C for 5 min, secondary denaturation at 95 °C for 1 min, annealing at 60 °C for 30 s, extension at 72 °C for 30 s, and final extension at 72 °C for 5 min. The standard strain of *K. pneumoniae* (ATCC BAA-1705) was used as a positive control. The amplified DNA fragment was then sent to the Gene Fanavaran Company for DNA sequencing.

**Antibiotic susceptibility test:** Disc diffusion method was used to measure the isolates antibiotic resistance pattern against 10 antibiotics according to the Clinical and Laboratory Standards Institute (CLSI) 2018 guidelines, including gentamicin (10 µg), streptomycin (10 µg), ampicillin (10 µg), meropenem (10 µg), ciprofloxacin (5 µg), nalidixic acid (30 µg), cotrimoxazole (1.25-23.75 µg), chloramphenicol (30 µg), ceftriaxone (30 µg), and levofloxacin (5 µg), prepared from Padtan Teb Company (Iran). For each clinical specimen, a standard opacity equivalent to 0.5 McFarland was prepared,

and after culturing the bacteria for 24 hours on Müeller-Hinton agar medium, antibiotic discs were placed on a plate and incubated for 24 hours at 37 °C. Then the diameter of the inhibition zone was measured with a ruler. Finally, antibiogram test report was recorded as susceptible, resistant, or semi-resistant. *K. pneumoniae* standard strain (ATCC BAA-1705) was used as the quality control.

**Hydroalcoholic extract preparation:**

About 20 g of dried and grounded leaves of *A. scoparias* (Herbarium number IBRC P1000352) were prepared from Iran Genetic Resources Center and extracted using percolation method [18].


**TiO<sub>2</sub> nanoparticles:** TiO<sub>2</sub> nanoparticles were purchased from the American company US Nano with an average particle size of 20 nm and a purity of 99.99%.

**MIC of *Artemisia scoparia* extract, TiO<sub>2</sub> nanoparticles, and the synergistic effect:**

Minimum inhibitory concentration method was used to determine the MIC of TiO<sub>2</sub> nanoparticles and *Artemisia* extract. The MIC method was performed using the microdilution method in a microplate according to the CLSI 2018 guideline and repeated 3 times. For this purpose, at the first step, 90 µL of Mueller Hinton broth medium was poured into the wells, and then 10 µL of microbial suspension at a concentration of 0.5 McFarland was added ( $1.5 \times 10^8$  CFU/mL bacterial concentration), then it was diluted at a ratio of 1:100 to obtain a density of  $10^6$  CFU/mL. After mixing with liquid solvents, the final concentration of bacteria was  $5 \times 10^5$  CFU/mL. In some wells, TiO<sub>2</sub> nanoparticles were poured at logarithmic concentrations of 2-32 µg/mL (5 wells for each isolate), and in some other wells, *Artemisia* extract was poured at logarithmic concentrations of 2-2048 µg/mL (11 wells for each isolate). In order to investigate their synergistic effect, logarithmic concentrations of nanoparticles and extract (5 wells for each

isolate) were used simultaneously. One well was considered as negative control (without bacteria), and one well was considered as positive control (without extract and nanoparticles). After 24 hours, the amount of light absorption was read at 600 nm using an ELISA reader (AWARENESS, TechnilogyINC, Atat fax 2100), and the concentration of antimicrobial agent in the first well with no visible bacterial growth was reported as its MIC values.

The MIC numbers were analyzed using SPSS software Version 26 by employing independent t-test. The significance level of the results between the groups (treatment with extract, nanoparticles, and simultaneous treatment) was evaluated with a  $p < .001$ .



**Figure 1)** Amplified fragment of the *hly* gene encoding hemolysin by PCR reaction. 1: *K. pneumoniae*, 2: 50 bp size marker.

## Findings

In this study, a total of 30 isolates of *K. pneumoniae* were studied and confirmed by phenotypic and genotypic methods. Then for final confirmation, the desired fragment of the *hly* gene was amplified by PCR and confirmed by comparing with a 50 bp DNA size marker. Figure 1 shows the result of electrophoresis of *hly* gene encoding hemolysin. The sequenced fragment was first checked by Chromas software, and then the sequence BLAST was performed on the NCBI site. The results indicated that the similarity of PCR product sequence with *hly* gene sequence in gene bank was more than 98%, indicating genetic confirmation of *K. pneumoniae* strain.


**Antibiotic resistance of isolates:** The antibiotic resistance of the isolates was evaluated by the disk diffusion method. The lowest resistance of the isolates was related to the levofloxacin (43%), and the highest antibiotic resistance was related to the streptomycin (81%) (Figure 2).

**MIC determination:** The results of MIC test demonstrated that the average MIC of the extract (896  $\mu\text{g/mL}$ ) was significantly ( $p < .001$ ) higher than that of TiO<sub>2</sub> nanoparticles (14.13  $\mu\text{g/mL}$ ), indicating the greater antimicrobial effect of TiO<sub>2</sub> nanoparticles. Also, simultaneous use of both antimicrobials exhibited a synergistic effect against 25 out of 30 isolates while their

**Table 1)** Comparing the MIC values of all the treatment groups using the independent t-test.

MIC ( $\mu\text{g/mL}$ )	Strain Number	Mean $\pm$ SE	Significance
<i>Artemisia scoparia</i> extract	30	896 $\pm$ 98	-
TiO <sub>2</sub>	30	14.13 $\pm$ 1.5 <sup>a</sup>	$P < .001$
<i>Artemisia scoparia</i> extract and TiO <sub>2</sub>	30	7.53 $\pm$ 0.78 <sup>b</sup>	$P < .001$

a: Compared MIC of the TiO<sub>2</sub> with *A. scoparia* extract, b: Compared MIC of simultaneous treatment with TiO<sub>2</sub> or *A. scoparia* extract.



**Figure 2)** *K. pneumoniae* isolates resistance pattern to selected antibiotics.

MIC values were reduced by one logarithmic concentration. The number of isolates treated by the synergistic effect (25 isolates) was also significantly higher ( $p < .001$ ) than the number of isolates not treated by the synergistic effect (5 isolates). Therefore, it was found that simultaneous use of *Artemisia* extract and  $TiO_2$  nanoparticles exerts a synergistic effect (Table 1).

### Discussion

Most *K. pneumoniae* infections are either hospital-acquired or observed in people with debilitating diseases. *K. pneumoniae* infection often occurs in hospitalized patients with impaired ciliary function and non-hospitalized patients with chronic obstructive pulmonary disease, diabetes, or alcoholism [19].

It has been reported that 30  $\mu g$  of  $TiO_2$  nanoparticles is able to produce a 9 mm inhibition zone diameter on *K. pneumoniae* strains using the disk diffusion method. This inhibition zone diameter for *Staphylococcus aureus* is 18 mm [20].

Verdier et al. (2014) indicated that  $TiO_2$  nanoparticles could show an antibacterial

effect within two hours [21]. Their study was conducted on *Escherichia coli* strains by employing two types of experiments. These researchers attributed the mechanism of antimicrobial action of  $TiO_2$  nanoparticles to their photocatalytic properties. In the present study, it was found that  $TiO_2$  nanoparticles with MIC values of 8-32  $\mu g/mL$  could exhibit antimicrobial properties.

Tambekar and Dahikar (2011) selected a number of plants native to India and tested their antibacterial properties against a number of bacteria, including *K. pneumoniae*. The results showed that most plants had antibacterial effect [22]. In the present study, it was also observed that *A. scoparia* extract exhibited antimicrobial properties on clinical isolates of *K. pneumoniae* at concentrations between 512 and 1024  $\mu g/mL$ .

Winnett et al. (2017) prepared 106 extracts from 40 native Australian plants and studied their antimicrobial effects on *K. pneumoniae* isolates [23]. According to the results, 81.1% of the extracts had the ability to inhibit the growth of this bacterium [16, 22]. Furthermore, in the current study, it was found that *A. scoparia* extract possesses antimicrobial

properties.

Gavanji and colleagues (2014) studied the antibacterial effect of three plant essential oils on *K. pneumoniae*, *Pseudomonas aeruginosa*, and *S. aureus*. The results indicated that all three essential oils had a significant effect on the studied bacteria and among them, *Zataria multiflora* essential oil showed a greater antibacterial effect, and *S. aureus* indicated the highest susceptibility [24]. They also reported the antibacterial effect of *Artemisia* plant against the studied bacteria, indicating that this family species own a strong antibacterial effect. Indeed, the strength point of the present study is that it was performed on clinical isolates, but standard strains were studied in Gavanji's study. However, the effective concentration of *Artemisia* essential oil in Gavanji's report was 100 µg, while in the present study, the average MIC of *Artemisia* hydroalcoholic extract was 896 µg/mL.

Haroun and Al-Kayali (2016) reported the synergistic effect of *Thymbra spicata* L. crude extract with antibiotics against MDR strains of *K. pneumoniae* and *S. aureus*. The fractional inhibitory concentration (FIC) index was measured between 0.25-2 for *K. pneumoniae* [25].

Hacioglu et al. (2017) studied the antimicrobial activity of 31 widely consumed herbal teas alone and in combination with antibiotics on the bacteria such as *P. aeruginosa*, *Acinetobacter baumannii*, *E. coli*, *K. pneumoniae*, *Enterococcus faecalis*, methicillin-resistant *S. aureus*, and *Candida albicans* yeast through the disc diffusion method [26]. None of the herbal teas showed significant bactericidal properties alone, but their simultaneous use with antibiotics increased their bactericidal properties. There was a significant synergistic property between herbal teas and ampicillin, ampicillin-sulbactam, and nystatin antibiotics. In the present study, it

was observed that *Artemisia* extract in the presence of TiO<sub>2</sub> nanoparticles exhibited a synergistic effect (with 7.53 µg/mL mean) against 25 out of 30 isolates of *K. pneumoniae*, this antimicrobial property was significant ( $p < .001$ ).

## Conclusion

Considering the importance of controlling infections caused by MDR strains of *K. pneumoniae* and their antibiotic resistance as well as the effect of simultaneous use of herbal compounds and nanoparticles on this bacterium to overcome antibiotic resistance, in this study, the synergistic effect of *Artemisia* extract and TiO<sub>2</sub> nanoparticles on *K. pneumoniae* isolates was studied. The obtained results indicated that this approach could be considered as a combination therapy for MDR strains of *K. pneumoniae* in the future.

**Acknowledgments:** This work was conducted by the private fund of an M.Sc thesis with 10130560962022 code. Authors would like to thank all the lab assistants of the Islamic Azad University, Central Tehran Branch.

**Ethical permission:** This work was approved by Ethical committee of Islamic Azad University.

**Conflicts of interests:** None declared by authors.

**Authors' contribution:** Conceptualization: MR, SH; Data curation and formal analysis: MR; Investigation: MR; Methodology and project administration: All authors; Supervision: MR; Validation: MR, SH; Writing of original draft: SS; Writing, reviewing, and editing: MR, SH.

**Fundings:** Private fund.

**Consent to participate:** A written informed was obtained from all patients.

## References

1. Adams-Sapper S, Nolen S, Donzelli GF, Lal M, Chen K, Justo da Silva LH, et al. Rapid

- induction of high-level carbapenem resistance in heteroresistant KPC-producing *Klebsiella pneumoniae*. *Antimicrob Agents Chemother.* 2015;59(6):3281-9.
2. Bengoechea JA, Sa Pessoa J. *Klebsiella pneumoniae* infection biology: Living to counteract host defences. *FEMS Microbiol Rev.* 2019;43(2):123-44.
  3. Karabinis A, Paramythiotou E, Mylona-Petropoulou D, Kalogeromitros A, Katsarelis N, Kontopidou F, et al. Colistin for *Klebsiella pneumoniae*-associated sepsis. *Clin Infect Dis.* 2004;38(1):e7-9.
  4. Soltan Dalal MM, Miremadi SA, Sharifi Yazdi MK, Rastegar Lari AA, Rajabi Z, Avadis YS. Antimicrobial resistance trends of *Klebsiella* spp. isolated from patient in Imam Khomeini hospital. *J Pavard Salamat.* 2012;6(4):21-7.
  5. Behzadian Nejad Q, Abdollahi A, Najar Peerayeh SH, Forouhesh Tehrani H. Evaluation of bla-ctx-mtype gene in multi drug resistance *Klebsiella pneumoniae* species isolated from clinical samples. *Razi J Med Sci.* 2008;15(60-61):37-45.
  6. Wyres KL, Lam MM, Holt KE. Population genomics of *Klebsiella pneumoniae*. *Nature Rev Microbiol.* 2020;18(6):344-59.
  7. Shakeel M, Jabeen F, Shabbir S, Asghar MS, Khan MS, Chaudhry AS. Toxicity of nano-titanium dioxide (TiO<sub>2</sub>-NP) through various routes of exposure: A Review. *Biol Trace Elem Res.* 1)172;2016):36-1.
  8. Shi H, Magaye R, Castranova V, Zhao J. Titanium dioxide nanoparticles: A review of current toxicological data. Part *Fibre Toxicol.* 2013;10(1):1-33.
  9. Chellappa M, Anjaneyulu U, Manivasagam G, Vijayalakshmi U. Preparation and evaluation of the cytotoxic nature of TiO<sub>2</sub> nanoparticles by direct contact method. *Int J Nanomed.* 10;2015(Suppl 1):31-41.
  10. Garcia-Contreras R, Scougall-Vilchis RJ, Contreras-Bulnes R, Ando Y, Kanda Y, Hibino Y, et al. Effects of TiO<sub>2</sub> nanoparticles on cytotoxic action of chemotherapeutic drugs against a human oral squamous cell carcinoma cell line. *In Vivo.* 2)28;2014):15-209.
  11. Khan ST, Ahmad J, Ahamed M, Jousset A. Sub-lethal doses of widespread nanoparticles promote antifungal activity in *Pseudomonas protegens* CHA0. *Sci Total Environ.* 2018;627:658-62.
  12. Nam SY, Han NR, Rah SY, Seo Y, Kim HM, Jeong HJ. Anti-inflammatory effects of *Artemisia scoparia* and its active constituent, 3,5-dicaffeoyl-epiquinic acid against activated mast cells. *Immunopharmacol Immunotoxicol.* 2018;40(1):52-8.
  13. Ramezani M, Fazli-Bazzaz BS, Saghafi-Khadem F, Dabaghian A. Antimicrobial activity of four *Artemisia* species of Iran. *Fitoterapia.* 2004;75(2):201-3.
  14. Sajid M, Rashid Khan MR, Shah NA, Waris TS, Younis T, Ullah S, et al. Evaluation of *Artemisia scoparia* for hemostasis promotion activity. *Pak J Pharm Sci.* 2017;30(5):1709-13.
  15. Cha JD, Jeong MR, Jeong SI, Moon SE, Kim JY, Kil BS, et al. Chemical composition and antimicrobial activity of the essential oils of *Artemisia scoparia* and *A. capillaris*. *Planta Med.* 2005;71(02):186-90.
  16. Singh HP, Mittal S, Kaur S, Batish DR, Kohli RK. Chemical composition and antioxidant activity of essential oil from residues of *Artemisia scoparia*. *Food Chem.* 2009;114(2):642-5.
  17. Candan ED, Aksöz N. *Klebsiella pneumoniae*: Characteristics of carbapenem resistance and virulence factors. *Acta Biochim Pol.* 2015;62(4).
  18. Singh J. Maceration, percolation, and infusion techniques for the extraction of medicinal and aromatic plants. *CIMAP.*

- 2008;67:32-5.
19. Martin RM, Bachman MA. Colonization, infection, and the accessory genome of *Klebsiella pneumoniae*. *Front Cell Infect Microbiol.* 2018;8:4.
  20. Ayati R, Dodi M, Karami M. Evaluation the effect of TiO<sub>2</sub> nanoparticles on MDR *Klebsiella pneumoniae* and *Staphylococcus aureus* strains resistant to multi antibiotics. 17<sup>th</sup> Iranian Biology conference 2012.
  21. Verdier T, Coutand M, Bertron A, Roques C. Antibacterial activity of TiO<sub>2</sub> photocatalyst alone or in coatings on *E. coli*: The influence of methodological Aspects. *Coatings.* 2014;4(3):670-86.
  22. Tambekar DH, Dahikar SB. Antibacterial activity of some Indian Ayurvedic preparations against enteric bacterial pathogens. *J Adv Pharm Technol Res.* 2011;2(1):24-9.
  23. Winnett V, Sirdaarta J, White A, Clarke FM, Cock IE, Inhibition of *Klebsiella pneumoniae* growth by selected Australian plants: Natural approaches for the prevention and management of ankylosing spondylitis. *Inflammopharmacology.* 2017;25(2):223-35.
  24. Gavanji S, Mohammadi E, Larki B, Bakhtari A. Antimicrobial and cytotoxic evaluation of some herbal essential oils in comparison with common antibiotics in bioassay condition. *Integr Med Res.* 2014;3(3):142-52.
  25. Haroun MF, Al-Kayali RS. Synergistic effect of *Thymbra spicata* L. extracts with antibiotics against multidrug-resistant *Staphylococcus aureus* and *Klebsiella pneumoniae* strains. *Iran J Basic Med Sci.* 2016;19(11):1193-200.
  26. Hacıoglu M, Dosler S, Birteksoz Tan AS, Otuk G. Antimicrobial activities of widely consumed herbal teas, alone or in combination with antibiotics: An in vitro study. *Peer J.* 2017;5:e3467.